

Ring of Polytopes, Quasi-symmetric functions and Fibonacci numbers

Victor M. Buchstaber

Nickolai Erokhovets.

Abstract

In this paper we study the ring $\mathcal{P}$ of combinatorial convex polytopes. We introduce the algebra of operators $\mathcal{D}$ generated by the operators d_k that send an n -dimensional polytope P^n to the sum of all its $(n-k)$ -dimensional faces. It turns out that $\mathcal{D}$ is isomorphic to the universal Leibnitz-Hopf algebra with the antipode $\chi(d_k) = (-1)^k d_k$. Using the operators d_k we build the generalized f -polynomial, which is a ring homomorphism from $\mathcal{P}$ to the ring $\text{Qsym}[t_1, t_2, \dots][\alpha]$ of quasi-symmetric functions with coefficients in $\mathbb{Z}[\alpha]$. The images of two polytopes coincide if and only if their flag f -vectors are equal. We describe the image of this homomorphism over the integers and prove that over the rationals it is a free polynomial algebra with dimension of the n -th graded component equal to the n -th Fibonacci number. This gives a representation of the Fibonacci series as an infinite product. The homomorphism is an isomorphism on the graded group BB generated by the polytopes introduced by Bayer and Billera to find the linear span of flag f -vectors of convex polytopes. This gives the group BB a structure of the ring isomorphic to $f(\mathcal{P})$. We show that the ring of polytopes has a natural Hopf comodule structure over the Rota-Hopf algebra of posets. As a corollary we build a ring homomorphism $l_\alpha: \mathcal{P} \rightarrow \mathcal{R}[\alpha]$ such that $F(l_\alpha(P)) = f(P)^*$, where F is the Ehrenborg quasi-symmetric function.

1 Introduction

Convex polytopes is a classical object of convex geometry. In recent times the solution of problems of convex geometry involves results of algebraic geometry and topology, commutative and homological algebra. There are remarkable results lying on the crossroads of the polytope theory, theory of complex manifolds, equivariant topology and singularity theory (see the survey [BR]). The example of successful collaboration of the polytope theory and differential equations can be found in [Buch].

To study the combinatorics of convex polytopes we develop the approach based on the ring $\mathcal{P}$ of combinatorial convex polytopes. In the focus of our interest we put the ring of linear operators $\mathcal{L}(\mathcal{P}) = \text{Hom}_{\mathbb{Z}}(\mathcal{P}, \mathcal{P})$, and its subring generated by the operators d_k , $k \geq 1$ that map an n -dimensional polytope to the sum of all its $(n-k)$ -dimensional faces. $d = d_1$ is a derivation, so $\mathcal{P}$ is a differential ring.

The ring $\mathcal{D}$ has a canonical structure of the Leibnitz-Hopf algebra (see Section 4). We prove that $\mathcal{D}$ is isomorphic to the universal Hopf algebra in the category of Leibnitz-Hopf algebras with the antipode $\chi(d_k) = (-1)^k d_k$.

$$\mathcal{D} \simeq \mathcal{Z}/J_{\mathcal{D}}$$

where $\mathcal{Z} = \mathbb{Z}\langle Z_1, Z_2, \dots \rangle$ is the universal Leibnitz-Hopf algebra – a free associative Hopf algebra with the comultiplication

$$\Delta Z_k = \sum_{i+j=k} Z_i \otimes Z_j,$$

and $J_{\mathcal{D}}$ is a two-sided Hopf ideal generated by the relations $\sum_{i+j=n} (-1)^i Z_i Z_j = 0$, $n \geq 1$.

This algebra appears in various application of theory of Hopf algebras in combinatorics: over the rationals it is isomorphic to the graded dual of the odd subalgebra $S_-(\text{Qsym}[t_1, t_2, \dots], \zeta_Q)$ [ABS, Remark 6.7], to the algebra of forms on chain operators [BLiu, Proposition 3.2], to the factor algebra of the algebra of Piery operators on a Eulerian poset by the ideal generated by the Euler relations [BMSW, Example 5.3].

The action of $\mathcal{D}$ on $\mathcal{P}$ satisfies the property $D_\omega(PQ) = \mu \circ (\Delta D_\omega)(P \otimes Q)$ so the ring $\mathcal{P}$ has the structure of a Milnor module over $\mathcal{D}$.

The graded dual Hopf algebra to $\mathcal{Z}$ is the algebra of quasi-symmetric functions $\text{Qsym}[t_1, t_2, \dots]$. The α -character $\xi_\alpha: \mathcal{P} \rightarrow \mathbb{Z}[\alpha]$ induces the homomorphism $\mathcal{P} \rightarrow \mathcal{D}^*[\alpha]$, and the composition with the inclusion $\mathcal{D}^* \subset \text{Qsym}[t_1, t_2, \dots]$ gives the ring homomorphism

$$f: \mathcal{P} \rightarrow \text{Qsym}[t_1, t_2, \dots][\alpha]$$

We call $f(\alpha, t_1, t_2, \dots)$ the generalized f -polynomial. It can be shown that $f(\alpha, t_1, 0, 0, \dots) = f_1(\alpha, t)$ is a homogeneous f -polynomial in two variables introduced in [Buch].

For simple polytopes $f(\alpha, t_1, t_2, \dots) = f_1(\alpha, t_1 + t_2 + \dots)$.

This paper has the following structure:

Sections 2-4 are devoted to the necessary definitions and important facts about quasi-symmetric functions and Hopf algebras.

Section 5 contains topological realizations of the Hopf algebras we study.

Section 6 recalls important facts about flag f -vectors and emphasizes some peculiarities we need in this paper.

Section 7 is devoted to the cone and the bipyramide operators C and B used by M. Bayer and L. Billera [BB] to find the linear span of flag f -vectors of polytopes. We show that the operations C and B can be defined on the ring $\text{Qsym}[t_1, t_2, \dots][\alpha]$ in such a way that $f(CP) = Cf(P)$ and $f(BP) = Bf(P)$.

In Section 8 the structure theorem for the ring $\mathcal{D}$ is proved.

In Section 9 we introduce the generalized f -polynomial and find the functional equations that describe the image of the ring $\mathcal{P}$ in $\text{Qsym}[t_1, t_2, \dots][\alpha]$. These equations are equivalent to the generalized Dehn-Sommerville equations discovered by M. Bayer and L. Billera in [BB].

In Section 10 the condition that determines the generalized f -polynomial in a unique way is found.

In Section 11 we describe the graded dual Hopf algebra to the Hopf algebra $\mathcal{D}$ as a subalgebra of $\text{Qsym}[t_1, t_2, \dots]$ and find the image of $\mathcal{P}$ in the ring $\mathcal{D}^*[\alpha]$.

In Section 12 the multiplicative structure of the ring $f(\mathcal{P} \otimes \mathbb{Q})$ is described. We prove that $f(\mathcal{P} \otimes \mathbb{Q})$ is a free polynomials algebra with dimension of n -th graded component equal to the n -th Fibonacci number c_n ($c_0 = c_1 = 1$, $c_{n+1} = c_n + c_{n-1}$). This gives the decomposition of the Fibonacci series into the infinite product

$$\frac{1}{1-t-t^2} = \sum_{n=0}^{\infty} c_n t^n = \prod_{i=1}^{\infty} \frac{1}{(1-t^i)^{k_i}},$$

where k_i is the number of generators of degree i . The infinite product converges absolutely in the interval $|t| < \frac{\sqrt{5}-1}{2}$. The numbers k_n satisfy the inequalities $k_{n+1} \geq k_n \geq N_n - 2$, where N_n is the number of the decompositions of n into the sum of odd numbers.

In Section 13 we introduce the multiplicative structure on the graded group BB generated by the Bayer-Billera polytopes arising from the isomorphism with the ring $f(\mathcal{P})$.

Nowadays Hopf algebras is one of the central tools in combinatorics. There is a well-known Hopf algebra of posets introduced by Joni and Rota in [JR]. Various aspects of this algebra were studied in [Ehr, ABS, Sch1, Sch2]. A generalization of the Rota-Hopf algebra was proposed in [RS]. In [Ehr] R. Ehrenborg introduced the F -quasi-symmetric function, which gives a Hopf algebra homomorphism from the Rota-Hopf algebra to $\text{Qsym}[t_1, t_2, \dots]$. In Section 14 we show that there is a natural right Hopf comodule structure over the Rota-Hopf algebra on the ring $\mathcal{P}$. It gives the natural ring homomorphism $l_\alpha: \mathcal{P} \rightarrow \mathcal{R}[\alpha]$. We prove that $F(l_\alpha(P^n)) = f(P^n)^*$ and show that the Hopf comodule structure agree with the homomorphisms f^* and F .

Section 15 deals with the problem of description of flag f -vectors of polytopes. For simple polytope g -theorem gives the full description of the set of flag f -vectors. As it is mentioned in [Z1] even for 4-dimensional non-simple polytopes the corresponding problem is extremely hard. We show how this problem can be stated in terms of the ring of polytopes.

2 The Ring of Polytopes

For the notion of a convex polytope see the books [Gb, Z2].

Definition 1. A *ring of polytopes* $\mathcal{P}$ is a ring of all combinatorial convex polytopes. As an abelian group $\mathcal{P} = \mathbb{Z} + \bigoplus_{n \geq 1, m-n \geq 1} \mathcal{P}^{n, m-n}$, where $\mathcal{P}^{n, m-n}$ is a free abelian group generated by all n -dimensional combinatorial polytopes with m facets. The direct product of polytopes $P \times Q$ defines a multiplication and the point $1 = \text{pt}$ is a unit.

It can be proved that $\mathcal{P}$ is a ring of polynomials generated by all indecomposable polytopes. n and $m - n$ are graduations of the ring and it easy to see that there is a finite number of n -dimensional polytopes with m facets, so an abelian group $\mathcal{P}^{n, m-n}$ is finitely generated.

Definition 2. Let us denote $\mathcal{P}^0 = \mathbb{Z}$, $\mathcal{P}^n = \bigoplus_{m=n+1}^{\infty} \mathcal{P}^{n, m-n}$, $n \geq 1$, and $\mathcal{P}^{[n]} = \bigoplus_{k=0}^n \mathcal{P}^k$.

We will use the notations: P for a polytope, P^n for a polytope of dimension n , p for an element of $\mathcal{P}$, p^n for an element of $\mathcal{P}^n$.

Definition 3. Let us define the ring of linear operators $\mathcal{L}(\mathcal{P}) = \text{Hom}_{\mathbb{Z}}(\mathcal{P}, \mathcal{P})$ with the multiplication given by a composition of operators.

Definition 4. Let d_k be an operator defined on each polytope P^n as the sum of all $(n - k)$ -dimensional faces.

$$d_k P^n = \sum_{F^{n-k} \subset P^n} F^{n-k}$$

For example, $d_0 = \text{id}$, and $d = d_1$ is a facets operator. Since $d(P \times Q) = (dP) \times Q + P \times (dQ)$, it is a derivation. So the ring of polytopes is a differential ring.

Let us denote by $\mathcal{D} = \mathcal{D}(\mathcal{P}) \subset \mathcal{L}(\mathcal{P})$ the subring of operators on $\mathcal{P}$ generated by d_k , $k \geq 1$.

The ring of polytopes contains a differential subring $\mathcal{P}_s$, generated by all simple polytopes.

In the ring of simple polytopes we have the relations: $d_k|_{\mathcal{P}_s} = \frac{d^k}{k!}|_{\mathcal{P}_s}$, so $\mathcal{D}(\mathcal{P}_s)$ is isomorphic to the divided power ring $d_i d_j = \binom{i+j}{i} d_{i+j}$.

In the general case as an abelian group the ring $\mathcal{D}$ is generated by the monomials $D_\omega = d_{j_1} \dots d_{j_k}$ where ω is a k -tuple $(j_1, \dots, j_k)$, for some $k \geq 0$, $j_i \geq 1$.

Definition 5. A *composition* ω of a number n is an ordered set $\omega = (j_1, \dots, j_k)$, $j_i \geq 1$, such that $n = j_1 + \dots + j_k$. Let us denote $|\omega| = n$.

The number of compositions of n into exactly k parts is given by the binomial coefficient $\binom{n-1}{k-1}$.

It is easy to see that

$$d_k(P \times Q) = \sum_{i+j=k} d_i P \times d_j Q = d_k P \times Q + d_{k-1} P \times dQ + \dots + P \times d_k Q.$$

and

$$D_\omega(P \times Q) = \sum_{\Omega' + \Omega'' = \omega} D_{\omega'} P \times D_{\omega''} Q. \quad (1)$$

where for the compositions $\omega = (j_1, \dots, j_k)$, $\omega' = (j'_1, \dots, j'_{l'})$, $\omega'' = (j''_1, \dots, j''_{l''})$, Ω' and Ω'' are all the k -tuples such that

$$\Omega' = (0, \dots, j'_1, \dots, 0, \dots, j'_{l'}, \dots, 0), \quad \Omega'' = (0, \dots, j''_1, \dots, 0, \dots, j''_{l''}, \dots, 0),$$

For example,

$$\begin{aligned} dd_3(P \times Q) &= (dd_3 P) \times (Q) + (dd_2 P) \times (dQ) + (d^2 P) \times (d_2 Q) + (dP) \times (d_3 Q) + \\ &\quad + (d_3 P) \times (dQ) + (d_2 P) \times (d^2 Q) + (dP) \times (dd_2 Q) + P \times (dd_3 Q); \end{aligned}$$

The list of summands corresponds to the decompositions:

$$\begin{aligned} (1, 3) &= (1, 3) + (0, 0) = (1, 2) + (0, 1) = (1, 1) + (0, 2) = (1, 0) + (0, 3) = \\ &= (0, 3) + (1, 0) = (0, 2) + (1, 1) = (0, 1) + (1, 2) = (0, 0) + (1, 3). \end{aligned}$$

This formula gives a coproduct in $\mathcal{D}$ defined on the generators by the formula

$$\Delta d_k = \sum_{i+j=k} d_i \otimes d_j$$

and thus $\mathcal{D}$ obtains the structure of a graded Hopf algebra with $\deg d_k = k$.

Definition 6 ([N]). A *Milnor module* M over the Hopf algebra X is an algebra with unit $1 \in k$ which is also a module over X satisfying

$$x(uv) = \sum x'_n(u)x''_n(v), \quad x \in X, u, v \in M, \quad \Delta x = \sum x'_n \otimes x''_n.$$

By definition of the comultiplication on the ring $\mathcal{D}$

$$\mu\Delta(D_\omega)(P \otimes Q) = D_\omega(PQ),$$

therefore for the ring $k = \mathbb{Z}$ we obtain the proposition:

Proposition 1. *The ring of polytopes $\mathcal{P}$ is a Milnor module over the Hopf algebra $\mathcal{D}$.*

Another example of operators in $\mathcal{L}(\mathcal{P})$ gives the multiplication by the elements of $\mathcal{P}$:

$$[P](Q) = PQ$$

Proposition 2. *The following relation holds*

$$d_k[P] = \sum_{i=0}^k [d_i P] d_{k-i}$$

Proof.

$$d_k(PQ) = \sum_{i=0}^k (d_i P) d_{k-i} Q$$

□

Thus any operator in the ring $\mathcal{PD}$ generated by $\mathcal{D}$ and $\{[P], P \in \mathcal{P}\}$ can be expressed as a sum of operators

$$\sum_{P, \omega} [P] D_\omega$$

Definition 7. Define an α -character $\xi_\alpha : \mathcal{P} \rightarrow \mathbb{Z}[\alpha]$ by the formula

$$\xi_\alpha P^n = \alpha^n.$$

For $\alpha = 0$ set

$$\xi_0(P^n) = \begin{cases} 1, & P^n = \text{pt}; \\ 0, & n \geq 1. \end{cases}$$

Let us denote $\varepsilon = \xi_0$.

Definition 8. Let us define an operator $\Phi : \mathcal{P} \rightarrow \mathcal{P}[t]$, $\Phi(t) \in \mathcal{D}[[t]]$:

$$\Phi(t) = 1 + dt + d_2 t^2 + d_3 t^3 + \cdots = \sum_{k=0}^{\infty} d_k t^k.$$

Proposition 3.

$$\xi_{-\alpha} \Phi(\alpha) = \xi_\alpha$$

Proof. Let us note that $\xi_\alpha d_k P^n = f_{n-k}(P^n)$ – the number of $(n-k)$ -dimensional faces. Then for any polytope P^n we have

$$\xi_{-\alpha} \Phi(\alpha) P^n = (-\alpha)^n + (-\alpha)^{n-1} f_{n-1} \alpha + \cdots + f_0 \alpha^n = \left(\sum_{i=0}^n (-1)^i f_i \right) \alpha^n = \alpha^n = \xi_\alpha P^n,$$

since the Euler formula gives the relation:

$$f_0 - f_1 + \cdots + (-1)^{n-1} f_{n-1} + (-1)^n = 1$$

□

Definition 9. Let P^n be the n -dimensional polytope

$$P^n = \{ \mathbf{x} \in \mathbb{R}^n : \langle \mathbf{a}_i, \mathbf{x} \rangle + 1 \geq 0 \}$$

Then a *dual* (or *polar*) *polytope* is defined as $P^* = \{ \mathbf{y} \in \mathbb{R}^n : \langle \mathbf{y}, \mathbf{x} \rangle + 1 \geq 0, \forall \mathbf{x} \in P^n \}$.

There is a bijection $F \longleftrightarrow F^\diamond$ between the i -faces of P and the $(n-1-i)$ -faces of P^* such that

$$F \subset G \Leftrightarrow G^\diamond \subset F^\diamond.$$

The operation $*$ defines a linear operator on the ring $\mathcal{P}$ that does not belong to $\mathcal{PD}$

3 Quasi-Symmetric Functions

Definition 10. Let $t_1, t_2, \dots$ be a finite or an infinite set of variables. For a composition $\omega = (j_1, \dots, j_k)$ consider a *quasi-symmetric monomial*

$$M_\omega = \sum_{l_1 < \dots < l_k} t_{l_1}^{j_1} \dots t_{l_k}^{j_k}.$$

Degree of the monomial M_ω is equal to $|\omega| = j_1 + \dots + j_k$.

For any two monomials $M_{\omega'}$ and $M_{\omega''}$ their product in the ring of polynomials $\mathbb{Z}[t_1, t_2, \dots]$ is equal to

$$M_{\omega'} M_{\omega''} = \sum_{\omega} \left(\sum_{\Omega' + \Omega'' = \omega} \right) M_\omega,$$

where for the compositions $\omega = (j_1, \dots, j_k)$, $\omega' = (j'_1, \dots, j'_{l'})$, $\omega'' = (j''_1, \dots, j''_{l''})$, Ω' and Ω'' are all the k -tuples such that

$$\Omega' = (0, \dots, j'_1, \dots, 0, \dots, j'_{l'}, \dots, 0), \quad \Omega'' = (0, \dots, j''_1, \dots, 0, \dots, j''_{l''}, \dots, 0),$$

This multiplication rule of compositions is called *the overlapping shuffle multiplication*.

For example,

1.

$$M_{(1)} M_{(1)} = \left(\sum_i t_i \right) \left(\sum_j t_j \right) = \sum_i t_i^2 + 2 \sum_{i < j} t_i t_j = M_{(2)} + 2M_{(1,1)}.$$

This corresponds to the decompositions

$$(2) = (1) + (1), \quad (1, 1) = (1, 0) + (0, 1) = (0, 1) + (1, 0).$$

2.

$$\begin{aligned} M_{(1)} M_{(1,1)} &= \left(\sum_i t_i \right) \left(\sum_{j < k} t_j t_k \right) = \sum_{i < j} t_i^2 t_j + \sum_{i < j} t_i t_j^2 + 3 \sum_{i < j < k} t_i t_j t_k = \\ &= M_{(2,1)} + M_{(1,2)} + 3M_{(1,1,1)}. \end{aligned}$$

This corresponds to the decompositions

$$(2, 1) = (1, 0) + (1, 1), \quad (1, 2) = (1, 0) + (1, 1), \\ (1, 1, 1) = (1, 0, 0) + (0, 1, 1) = (0, 1, 0) + (1, 0, 1) = (0, 0, 1) + (1, 1, 0).$$

3.

$$\begin{aligned} M_{(1,1)}M_{(1,1)} &= \left(\sum_{i < j} t_i t_j \right) \left(\sum_{k < l} t_k t_l \right) = \sum_{i < j} t_i^2 t_j^2 + 2 \sum_{i < j < k} t_i^2 t_j t_k + \\ &\quad + 2 \sum_{i < j < k} t_i t_j^2 t_k + 2 \sum_{i < j < k} t_i t_j t_k^2 + 6 \sum_{i < j < k < l} t_i t_j t_k t_l = \\ &= M_{(2,2)} + 2M_{(2,1,1)} + 2M_{(1,2,1)} + 2M_{(1,1,2)} + 6M_{(1,1,1,1)}. \end{aligned}$$

This corresponds to the decompositions

$$\begin{aligned} (2, 2) &= (1, 1) + (1, 1), \quad (2, 1, 1) = (1, 1, 0) + (1, 0, 1) = (1, 0, 1) + (1, 1, 0), \\ (1, 2, 1) &= (1, 1, 0) + (0, 1, 1) = (0, 1, 1) + (1, 1, 0), \quad (1, 1, 2) = (1, 0, 1) + (0, 1, 1) = (0, 1, 1) + (1, 0, 1), \\ (1, 1, 1, 1) &= (1, 1, 0, 0) + (0, 0, 1, 1) = (1, 0, 1, 0) + (0, 1, 0, 1) = (1, 0, 0, 1) + (0, 1, 1, 0) = \\ &= (0, 1, 1, 0) + (1, 0, 0, 1) = (0, 1, 0, 1) + (1, 0, 1, 0) = (0, 0, 1, 1) + (1, 1, 0, 0). \end{aligned}$$

Thus finite integer combinations of quasi-symmetric monomials form a ring. This ring is called *a ring of quasi-symmetric functions* and is denoted by $\text{Qsym}[t_1, \dots, t_n]$ (where n is the number of variables. In the case of an infinite number of variables it is denoted by $\text{Qsym}[t_1, t_2, \dots]$).

Proposition 4. *A polynomial $g \in \mathbb{Z}[t_1, \dots, t_m]$ is a finite linear combination of quasi-symmetric monomials if and only if*

$$g(0, t_1, t_2, \dots, t_{m-1}) = g(t_1, 0, t_2, \dots, t_{m-1}) = \dots = g(t_1, \dots, t_{m-1}, 0)$$

Proof. For the quasi-symmetric monomial M_ω we have

$$M_\omega(t_1, \dots, t_i, 0, t_{i+1}, \dots, t_{m-1}) = \begin{cases} 0, & \omega = (j_1, \dots, j_m); \\ M_\omega(t_1, \dots, t_i, t_{i+1}, \dots, t_{m-1}), & \omega = (j_1, \dots, j_k), \quad k < m. \end{cases}$$

So this property is true for all quasi-symmetric functions.

On the other hand, let the condition of the proposition be true. Let us prove that for a fixed composition ω any two monomials $t_{l_1}^{j_1} \dots t_{l_k}^{j_k}$ and $t_{l'_1}^{j_1} \dots t_{l'_k}^{j_k}$ have the equal coefficients $g_{l_1, \dots, l_k}^{j_1, \dots, j_k}$ and $g_{l'_1, \dots, l'_k}^{j_1, \dots, j_k}$.

Let $l_i + 1 < l_{i+1}$ or $i = k$ and $l_i < m$. Consider the corresponding coefficients in the polynomial equation:

$$g(t_1, \dots, t_{l_i-1}, 0, t_{l_i}, t_{l_i+1}, \dots, t_{m-1}) = g(t_1, \dots, t_{l_i-1}, t_{l_i}, 0, t_{l_i+1}, \dots, t_{m-1}).$$

On the left the monomial $t_{l_1}^{j_1} \dots t_{l_i}^{j_i} t_{l_{i+1}-1}^{j_{i+1}} \dots t_{l_k}^{j_k}$ has the coefficient $g_{l_1, \dots, l_i, l_{i+1}, l_{i+1}, \dots, l_k}^{j_1, \dots, j_k}$ and on the right $g_{l_1, \dots, l_i, l_i, l_{i+1}, \dots, l_k}^{j_1, \dots, j_k}$, so they are equal. Now we can move the index l_i to the right to $l_i + 1$ and in the same manner we can move l_i to the left, if $l_{i-1} < l_i - 1$ or $i = 1$ and $l_i > 1$.

Now let us move step by step the index l_1 to 1, then the index l_2 to 2, and so on. At last we obtain that $g_{l_1, \dots, l_k}^{j_1, \dots, j_k} = g_{1, \dots, k}^{j_1, \dots, j_k}$. $\square$

The same is true in the case of an infinite number of variables, if we consider all the expressions

$$r + \sum_{k=1}^{\infty} \sum_{1 \leq l_1 < \dots < l_k} \sum_{j_1, \dots, j_k \geq 1} r_{l_1, \dots, l_k}^{j_1, \dots, j_k} t_{l_1}^{j_1} \dots t_{l_k}^{j_k}.$$

of bounded degree: $|\omega| = j_1 + \dots + j_k < N$. As a corollary we obtain another proof of the fact that quasi-symmetric functions form a ring.

In [H] M. Hazewinkel proved the Ditters conjecture that $\text{Qsym}[t_1, t_2, \dots]$ is a free commutative algebra of polynomials over the integers.

The n -th graded component $\text{Qsym}^n[t_1, t_2, \dots]$ has rank 2^{n-1} .

The numbers β_i of the multiplicative generators of degree i can be found by a recursive relation:

$$\frac{1-t}{1-2t} = \prod_{i=1}^{\infty} \frac{1}{(1-t^i)^{\beta_i}}$$

There are ring homomorphisms:

$$\begin{aligned} v_{m+1} : \text{Qsym}[t_1, \dots, t_m] &\rightarrow \text{Qsym}[t_1, \dots, t_{m+1}] : & v_{m+1} M_\omega(t_1, \dots, t_m) &= M_\omega(t_1, \dots, t_{m+1}) \\ u_m : \text{Qsym}[t_1, \dots, t_{m+1}] &\rightarrow \text{Qsym}[t_1, \dots, t_m] : & t_{m+1} &\rightarrow 0; \end{aligned}$$

The mapping u_m sends all the monomials M_ω corresponding to the compositions $(j_1, \dots, j_{m+1})$ of length $m+1$ to zero, and for the compositions ω of smaller length $u_m M_\omega(t_1, \dots, t_{m+1}) = M_\omega(t_1, \dots, t_m)$.

It is easy to see that $u_m v_{m+1}$ is the identity map of the ring $\text{Qsym}[t_1, \dots, t_m]$.

Given $m > 0$ there is a projection $\Pi_{\text{Qsym}} : \mathbb{Z}[t_1, \dots, t_m] \rightarrow \text{Qsym}[t_1, \dots, t_m] \otimes \mathbb{Q}$:
for $j_1 < j_2 < \dots < j_k$

$$\Pi_{\text{Qsym}} t_{j_1}^{j_1} \dots t_{j_k}^{j_k} = \frac{1}{\binom{m}{k}} \left(\sum_{l_1 < \dots < l_k} t_{l_1}^{j_1} \dots t_{l_k}^{j_k} \right),$$

which gives the average value over all the monomials of each type.

Then for the quasi-symmetric monomial M_ω , $\omega = (j_1, \dots, j_k)$ we have:

$$\Pi_{\text{Qsym}} M_\omega = \sum_{1 \leq j_1 < \dots < j_k \leq m} \frac{1}{\binom{m}{k}} M_\omega = \binom{m}{k} \frac{1}{\binom{m}{k}} M_\omega = M_\omega,$$

So Π_{Qsym} is indeed a projection.

Remark. We see that in the theory of symmetric and quasi-symmetric functions there is an important additional graduation – the number of variables in the polynomial.

Definition 11. For a composition $\omega = (j_1, \dots, j_k)$ let us define the composition $\omega^* = (j_k, \dots, j_1)$.

The correspondence $M_\omega \rightarrow (M_\omega)^* = M_{\omega^*}$ defines an involutory ring homomorphism

$$* : \text{Qsym}[t_1, t_2, \dots] \rightarrow \text{Qsym}[t_1, t_2, \dots].$$

4 Hopf Algebras

In this part we follow mainly the notations of [H]. See also [BR] and [CFL].

Let R be a commutative associative ring with unity.

4.1 Leibnitz-Hopf Algebras

Definition 12. A *Leibnitz-Hopf algebra* over the ring R is an associative Hopf algebra $\mathcal{H}$ over the ring R with a fixed sequence of a finite or countable number of multiplicative generators H_i , $i = 1, 2, \dots$ satisfying the comultiplication formula

$$\Delta H_n = \sum_{i+j=n} H_i \otimes H_j, \quad H_0 = 1.$$

A *universal Leibnitz-Hopf algebra* $\mathcal{A}$ over the ring R is a Leibnitz-Hopf algebra with the universal property: for any Leibnitz-Hopf algebra $\mathcal{H}$ over the ring R there exists a unique Hopf algebra homomorphism $\mathcal{A} \rightarrow \mathcal{H}$.

Consider the free associative Leibnitz-Hopf algebra over the integers $\mathcal{Z} = \mathbb{Z}\langle Z_1, Z_2, \dots \rangle$ in countably many generators Z_i .

Proposition 5. $\mathcal{Z}$ is a universal Leibnitz-Hopf algebra over the integers, and $\mathcal{Z} \otimes R$ is a universal Leibnitz-Hopf algebra over the ring R .

Set $\deg Z_i = i$. Let us denote by $\mathcal{M}$ the graded dual Hopf algebra over the integers.

It is not difficult to see that $\mathcal{M}$ is precisely the algebra of quasi-symmetric functions over the integers. Indeed, for any composition $\omega = (j_1, \dots, j_k)$ we can define m_ω by the dual basis formula

$$\langle m_\omega, Z^\sigma \rangle = \delta_{\omega, \sigma},$$

where $Z^\sigma = Z_{r_1} \dots Z_{r_l}$ for a composition $\sigma = (r_1, \dots, r_l)$. Then the elements m_ω multiply exactly as the quasi-symmetric monomials M_ω .

Let us denote

$$\Phi(t) = 1 + Z_1 t + Z_2 t^2 + \dots = \sum_{k=0}^{\infty} Z_k t^k$$

Then the comultiplication formula is equivalent to

$$\Delta \Phi(t) = \Phi(t) \otimes \Phi(t)$$

Let $\chi : \mathcal{Z} \rightarrow \mathcal{Z}$ be the *antipode*, that is a linear operator, satisfying the property

$$1 \star \chi = \mu \circ 1 \otimes \chi \circ \Delta = \eta \circ \varepsilon = \mu \circ \chi \otimes 1 \circ \Delta = \chi \star 1,$$

where $\varepsilon : \mathcal{Z} \rightarrow \mathbb{Z}$, $\varepsilon(1) = 1$, $\varepsilon(Z^\sigma) = 0$, $\sigma \neq \emptyset$, is a *counit*, $\eta : \mathbb{Z} \rightarrow \mathcal{Z}$, $\eta(a) = a \cdot 1$ is a *unite* map, and μ is a multiplication in $\mathcal{Z}$.

Then $\Phi(t)\chi(\Phi(t)) = 1 = \chi(\Phi(t))\Phi(t)$ and $\{\chi(Z_n)\}$ satisfy the recurrent formulas

$$\chi(Z_1) = -Z_1; \quad \chi(Z_{n+1}) + \chi(Z_n)Z_1 + \dots + \chi(Z_1)Z_n + Z_{n+1} = 0, \quad n \geq 1. \quad (2)$$

In fact, since $\Phi(t) = 1 + Z_1 t + \dots$, we can use the previous formula to obtain

$$\begin{aligned} \chi(\Phi(t)) &= \frac{1}{\Phi(t)} = \sum_{i=0}^{\infty} (-1)^i (\Phi(t) - 1)^i; \\ \chi(Z_n) &= \sum_{m=1}^n (-1)^m \sum_{a_1 + \dots + a_m = n} d_{a_1} \dots d_{a_m} \quad a_i \geq 1, \end{aligned}$$

This formula defines χ on generators, and therefore on the whole algebra.

The homomorphism $R : \mathcal{Z} \rightarrow \mathcal{D} : Z_k \rightarrow d_k$, defines in the ring of polytopes the structure of a right module over the Leibnitz-Hops algebra $\mathcal{Z}$. Let us mention, that the word $Z^\omega = Z_{j_1} \dots Z_{j_k}$ corresponds to the operator $D_{\omega^*} = d_{j_k} d_{j_{k-1}} \dots d_{j_1}$.

Definition 13. A *universal commutative Leibnitz-Hopf algebra* $\mathcal{C} = \mathbb{Z}[C_1, C_2, \dots]$ is a free commutative polynomial Leibnitz-Hopf algebra in generators C_i of degree i . We have $\mathcal{C} = \mathcal{Z}/J_{\mathcal{C}}$, where the ideal $J_{\mathcal{C}}$ is generated by the relations $Z_i Z_j - Z_j Z_i$.

It is a self-dual Hopf algebra and the graded dual Hopf algebra is naturally isomorphic to the algebra of symmetric functions $\mathbb{Z}[\sigma_1, \sigma_2, \dots] = \text{Sym}[t_1, t_2, \dots] \subset \text{Qsym}[t_1, t_2, \dots]$ generated by the symmetric monomials

$$\sigma_i = M_{\omega_i} = \sum_{l_1, \dots, l_i} t_{l_1} \dots t_{l_i},$$

where $\omega_i = (\underbrace{1, \dots, 1}_i)$.

The isomorphism $\mathcal{C} \simeq \mathcal{C}^*$ is given by the correspondence $C_i \rightarrow \sigma_i$.

4.2 Lie-Hopf Algebras

Definition 14. A *Lie-Hopf algebra* over the ring R is an associative Hopf algebra $\mathcal{L}$ with a fixed sequence of a finite or countable number of multiplicative generators $L_i, i = 1, 2, \dots$ satisfying the comultiplication formula

$$\Delta L_i = 1 \otimes L_i + L_i \otimes 1, \quad L_0 = 1.$$

A *universal Lie-Hopf algebra* over the ring R is a Lie-Hopf algebra $\mathcal{A}$ satisfying the universal property: for any Lie-Hopf algebra $\mathcal{H}$ over the ring R there exists a unique Hopf algebra homomorphism $\mathcal{A} \rightarrow \mathcal{H}$.

Consider the free associative Lie-Hopf algebra over the integers $\mathbb{Z}\langle U_1, U_2, \dots \rangle$ in countably many variables U_i .

Proposition 6. $\mathcal{U}$ is a universal Lie-Hopf algebra, and $\mathcal{U} \otimes R$ is a universal Lie-Hopf algebra over the ring R .

Let us set $\deg U_i = i$. Then the graded dual Hopf algebra is denoted by $\mathcal{N}$. This is the so-called *shuffle algebra*.

The antipode χ in $\mathcal{U}$ has a very simple form $\chi(U_i) = -U_i$.

4.3 Lyndon Words

A well-known theorem in the theory of free Lie algebras (see, for example, [Re]) states that the algebra $\mathcal{N} \otimes \mathbb{Q}$ is a commutative free polynomial algebra in the so-called *Lyndon words*.

Definition 15. Let us denote by $[a_1, \dots, a_n]$ an element of $\mathbb{N}^*$, that is the word over $\mathbb{N}$, consisting of symbols $a_1, \dots, a_n, a_i \in \mathbb{N}$. Let us order the words from $\mathbb{N}^*$ lexicographically, where any symbol is larger than nothing, that is $[a_1, \dots, a_n] > [b_1, \dots, b_m]$ if and only if there is an i such that $a_1 = b_1, \dots, a_{i-1} = b_{i-1}, a_i > b_i, 1 \leq i \leq \min\{m, n\}$, or $n > m$ and $a_1 = b_1, \dots, a_m = b_m$.

A *proper tail* of a word $[a_1, \dots, a_n]$ is a word of the form $[a_i, \dots, a_n]$ with $1 < i \leq n$. (The empty word and one-symbol word have no proper tails.)

A word is *Lyndon* if all its proper tails are larger than the word itself. For example, the words $[1, 1, 2], [1, 2, 1, 2, 2], [1, 3, 1, 5]$ are Lyndon and the words $[1, 1, 1, 1], [1, 2, 1, 2], [2, 1]$ are not Lyndon. The set of Lyndon words is denoted by LYN.

The same definitions make sense for any totally ordered set, for example, for the set $\{1, 2\}$ or for the set of all odd positive integers.

The role of Lyndon words is described by the following theorem:

Theorem (Chen-Fox-Lyndon Factorization [CFL]). *Every word w in $\mathbb{N}^*$ factors uniquely into a decreasing concatenation product of Lyndon words*

$$w = v_1 * v_2 * \dots * v_k, \quad v_i \in \text{LYN}, \quad v_1 \geq v_2 \geq \dots \geq v_k$$

For example, $[1, 1, 1, 1] = [1] * [1] * [1] * [1]$, $[1, 2, 1, 2] = [1, 2] * [1, 2]$, $[2, 1] = [2] * [1]$.

The algebra $\mathcal{N}$ is additively generated by the words in $\mathbb{N}^*$. The word $w = [a_1, \dots, a_n]$ corresponds to the function

$$\langle [a_1, \dots, a_n], U^\sigma \rangle = \delta_{w, \sigma}$$

where σ is a composition $\sigma = (r_1, \dots, r_l)$, $U^\sigma = U_{r_1} \dots U_{r_l}$, and for the word $[a_1, \dots, a_n]$

$$\delta_{w, \sigma} = \begin{cases} 1, & \sigma = (a_1, \dots, a_n); \\ 0, & \text{else} \end{cases}$$

The multiplication in the algebra $\mathcal{N}$ is the so-called *shuffle multiplication*:

$$[a_1, \dots, a_n] \times_{sh} [a_{n+1}, \dots, a_{m+n}] = \sum_{\sigma} [a_{\sigma(1)}, \dots, a_{\sigma(n)}, a_{\sigma(n+1)}, \dots, a_{\sigma(n+m)}],$$

where σ runs over all the substitutions $\sigma \in S_{m+n}$ such that

$$\sigma^{-1}(1) < \dots < \sigma^{-1}(n) \text{ and } \sigma^{-1}(n+1) < \dots < \sigma^{-1}(n+m).$$

For example,

$$\begin{aligned} [1] \times_{sh} [1] &= [1, 1] + [1, 1] = 2[1, 1]; \\ [1] \times_{sh} [2, 3] &= [1, 2, 3] + [2, 1, 3] + [2, 3, 1]; \\ [1, 2] \times_{sh} [1, 2] &= [1, 2, 1, 2] + [1, 1, 2, 2] + [1, 1, 2, 2] + [1, 1, 2, 2] + [1, 1, 2, 2] + [1, 2, 1, 2] = 2[1, 2, 1, 2] + 4[1, 1, 2, 2]. \end{aligned}$$

There is a well-known shuffle algebra structure theorem:

Theorem. $\mathcal{N} \otimes \mathbb{Q} = \mathbb{Q}[\text{LYN}]$, the free commutative algebra over $\mathbb{Q}$ in the symbols from LYN.

The proof follows from the following theorem concerning shuffle products in connection with Chen-Fox-Lyndon factorization.

Theorem. Let w be a word on the natural numbers and let $w = v_1 * v_2 * \dots * v_m$ be its Chen-Fox-Lyndon factorization. Then all words that occur with nonzero coefficient in the shuffle product $v_1 \times_{sh} v_2 \times_{sh} \dots \times_{sh} v_m$ are lexicographically smaller or equal to w , and w occurs with a nonzero coefficient in this product.

Given this result it is easy to prove the shuffle algebra structure theorem in the case of arbitrary totally ordered subset $M = \{m_1, m_2, \dots\} \subset \mathbb{N}$. Let us denote by $\mathcal{U}_M$ the free associative Lie-Hopf algebra $\mathbb{Z}\langle U_{m_1}, U_{m_2}, \dots \rangle$, let $\mathcal{N}_M$ be its graded dual algebra, and LYN_M be the corresponding set of all Lyndon words. We need to prove that $\mathcal{N}_M \otimes \mathbb{Q} = \mathbb{Q}[\text{LYN}_M]$.

Let m_1 be the minimal number in M . The smallest word $[m_1]$ is Lyndon.

Given a word w we can assume by induction that all the words lexicographically smaller than w have been written as polynomials in the elements of LYN_M . Take the Chen-Fox-Lyndon factorization $w = v_1 * v_2 * \dots * v_m$ of w and consider, using the preceding theorem,

$$v_1 \times_{sh} v_2 \times_{sh} \dots \times_{sh} v_m = aw + (\text{remainder}).$$

By the theorem the coefficient a is nonzero and all the words in (remainder) are lexicographically smaller than w , hence they belong to $\mathbb{Q}[\text{LYN}_M]$. Therefore $w \in \mathbb{Q}[\text{LYN}_M]$. This proves generation. Since each monomial in n -th graded component of $\mathcal{N} \otimes \mathbb{Q}$ has a unique Chen-Fox-Lyndon decomposition, the number of monomials in $\mathbb{Q}[\text{LYN}_M]$ of graduation n is equal to the number of monomials in $\mathcal{N}_M \otimes \mathbb{Q}$, monomials in Lyndon words are linearly independent. This proves that Lyndon words are algebraically independent.

Corollary 7. The graded dual algebras to the free associative Lie-Hopf algebras $\mathcal{U}_{12} = \mathbb{Z}\langle U_1, U_2 \rangle$ and $\mathcal{U}_{odd} = \mathbb{Z}\langle U_1, U_3, U_5, \dots \rangle$ are free polynomial algebras in Lyndon words LYN_{12} and LYN_{odd} respectively.

The correspondence

$$1 + Z_1 t + Z_2 t^2 + \dots = \exp(U_1 t + U_2 t^2 + \dots);$$

defines an isomorphism of Hopf algebras $\mathcal{Z} \otimes \mathbb{Q} \simeq \mathcal{U} \otimes \mathbb{Q}$.

However it is not true that $\mathcal{N}$ is a free polynomial commutative algebra over the integers.

5 Topological realization of Hopf algebras

In [BR] A. Baker and B. Richter showed that the ring of quasi-symmetric functions has a very nice topological interpretation.

We will consider CW -complexes X and their homology $H_*(X)$ and cohomology $H^*(X)$ with integer coefficients.

Let us assume that homology groups $H_*(X)$ have no torsion.

The diagonal map $X \rightarrow X \times X$ defines in $H_*(X)$ the structure of a graded coalgebra with the comultiplication $\Delta: H_*(X) \rightarrow H_*(X) \otimes H_*(X)$ and the dual structure of a graded algebra in $H^*(X)$ with the multiplication $\Delta^*: H^*(X) \otimes H^*(X) \rightarrow H^*(X)$.

In the case when X is an H -space with the multiplication $\mu: X \times X \rightarrow X$ we obtain the Pontryagin product $\mu_*: H_*(X) \otimes H_*(X) \rightarrow H_*(X)$ in the coalgebra $H_*(X)$ and the corresponding structure of a

graded Hopf algebra on $H_*(X)$. The cohomology ring $H^*(X)$ obtains the structure of a graded dual Hopf algebra with the diagonal mapping $\mu^*: H^*(X) \rightarrow H^*(X) \otimes H^*(X)$.

For any space Y the loop space $X = \Omega Y$ is an H -space. A continuous mapping $f: Y_1 \rightarrow Y_2$ induces a mapping of H -spaces $\Omega f: X_1 \rightarrow X_2$, where $X_i = \Omega Y_i$. Thus for any space Y such that $X = \Omega Y$ has no torsion in integral homology we obtain the Hopf algebra $H_*(X)$. This correspondence is functorial, that is any continuous mapping $f: Y_1 \rightarrow Y_2$ induces a Hopf algebra homomorphism $f_*: H_*(X_1) \rightarrow H_*(X_2)$.

By the Bott-Samelson theorem [BS], $H_*(\Omega \Sigma X)$ is the free associative algebra $T(\tilde{H}_*(X))$ generated by $\tilde{H}_*(X)$. This construction is functorial, that is a continuous mapping $f: X_1 \rightarrow X_2$ induces a ring homomorphism of the corresponding tensor algebras arising from the mapping $f_*: \tilde{H}_*(X_1) \rightarrow \tilde{H}_*(X_2)$. Denote elements of $H_*(\Omega \Sigma X)$ by $(a_1 | \dots | a_n)$, where $a_i \in \tilde{H}_*(X)$. Since the diagonal mapping $\Sigma X \rightarrow \Sigma X \times \Sigma X$ gives the H -map $\Delta: \Omega \Sigma X \rightarrow \Omega \Sigma X \times \Omega \Sigma X$, it follows from the Eilenberg-Zilber theorem that

$$\Delta_*(a_1 | \dots | a_n) = (\Delta_* a_1 | \dots | \Delta_* a_n)$$

where $(a_1 \otimes b_1 | a_2 \otimes b_2) = (a_1 | a_2) \otimes (b_1 | b_2)$.

There is a nice combinatorial model for any topological space of the form $\Omega \Sigma X$ with X connected, namely the James construction JX on X . After one suspension this gives rise to a splitting

$$\Sigma \Omega \Sigma X \sim \Sigma JX \sim \bigvee_{n \geq 1} \Sigma X^{(n)},$$

where $X^{(n)}$ denotes the n -fold smash power of X .

Example. There exists a homotopy equivalence

$$\Sigma \Omega \Sigma S^2 \rightarrow \bigvee_{n \geq 1} \Sigma (S^2)^{(n)} \simeq \Sigma \left(\bigvee_{n \geq 1} S^{2n} \right).$$

Therefore there exists an H -map $\Omega \Sigma (\Omega \Sigma S^2) \rightarrow \Omega \Sigma \left(\bigvee_{n \geq 1} S^{2n} \right)$ that is a homotopy equivalence.

Using these classical topological results let us describe topological realizations of the Hopf algebras we study in this work.

We have the following Hopf algebras:

- I. 1. $H_*(\mathbb{C}P^\infty)$ is a divided power algebra $\mathbb{Z}[v_1, v_2, \dots]/I$, where the ideal I is generated by the relations $v_n v_m - \binom{n+m}{n} v_{n+m}$, with the comultiplication

$$\Delta v_n = \sum_{k=0}^n v_k \otimes v_{n-k}; \quad (3)$$

2. $H^*(\mathbb{C}P^\infty) = \mathbb{Z}[u]$ – a polynomial ring with the comultiplication

$$\Delta u = 1 \otimes u + u \otimes 1;$$

- II. 1. $H_*(\Omega \Sigma \mathbb{C}P^\infty) \simeq T(\tilde{H}_*(\mathbb{C}P^\infty)) = \mathbb{Z}\langle v_1, v_2, \dots \rangle$ with v_i being non-commuting variables of degree $2i$. Thus there is an isomorphism of Hopf algebras

$$H_*(\Omega \Sigma \mathbb{C}P^\infty) \simeq \mathcal{Z}$$

under which v_n corresponds to Z_n .

The coproduct Δ on $H_*(\Omega \Sigma \mathbb{C}P^\infty)$ induced by the diagonal in $\Omega \Sigma \mathbb{C}P^\infty$ is compatible with the one in $\mathcal{Z}$:

$$\Delta v_n = \sum_{i+j=n} v_i \otimes v_j$$

So if we set $\deg Z_i = 2i$, then there is an isomorphism of graded Hopf algebras. This gives a geometric interpretation for the antipode χ in $\mathcal{Z}$: in $H_*(\Omega \Sigma \mathbb{C}P^\infty)$ it arises from the time-inversion of loops.

2. $H^*(\Omega\Sigma\mathbb{C}P^\infty)$ is the graded dual Hopf algebra to $H_*(\Omega\Sigma\mathbb{C}P^\infty)$.

Theorem. [BR] Let us double the graduation, that is let us set $\deg Z_i = 2i$, $\deg t_j = 2j$. Then there is an isomorphism of graded Hopf algebras:

$$H_*(\Omega\Sigma\mathbb{C}P^\infty) \simeq \mathcal{Z} = \mathbb{Z}\langle Z_1, Z_2, \dots \rangle, \quad H^*(\Omega\Sigma\mathbb{C}P^\infty) \simeq \mathcal{M} = \text{Qsym}[t_1, t_2, \dots].$$

III. $H_*(BU) \simeq H^*(BU) \simeq \mathbb{Z}[\sigma_1, \sigma_2, \dots] \simeq \mathcal{C}$. It is a self-dual Hopf algebra of symmetric functions. In the cohomology σ_i are represented by Chern classes.

IV. 1. $H_*(\Omega\Sigma S^2) = H_*(\Omega S^3) = \mathbb{Z}[v]$ – a polynomial ring with $\deg v = 2$ and the comultiplication

$$\Delta v = 1 \otimes v + v \otimes 1;$$

2. $H^*(\Omega\Sigma S^2) = \mathbb{Z}[v_n]/I$ is a divided power algebra. Thus $H^*(\Omega\Sigma S^2) \simeq H_*(\mathbb{C}P^\infty)$.

V. $H_*(\Omega\Sigma(\Omega\Sigma S^2)) \simeq \mathbb{Z}\langle w_1, w_2, \dots \rangle$. It is a free associative Hopf algebra with the comultiplication

$$\Delta w_n = \sum_{k=0}^n \binom{n}{k} w_k \otimes w_{n-k} \quad (4)$$

VI. $H_*\left(\Omega\Sigma\left(\bigvee_{n \geq 1} S^{2n}\right)\right) \simeq \mathbb{Z}\langle \xi_1, \xi_2, \dots \rangle$. It is a free associative algebra and has the structure of a graded Hopf algebra with the comultiplication

$$\Delta \xi_n = 1 \otimes \xi_n + \xi_n \otimes 1. \quad (5)$$

Therefore, $H_*\left(\Omega\Sigma\left(\bigvee_{n \geq 1} S^{2n}\right)\right)$ gives a topological realization of the universal Lie-Hopf algebra $\mathcal{U}$.

The homotopy equivalence

$$a: \Omega\Sigma\left(\bigvee_{n \geq 1} S^{2n}\right) \longrightarrow \Omega\Sigma(\Omega\Sigma S^2)$$

induces an isomorphism of graded Hopf algebras

$$a_*: \mathbb{Z}\langle \xi_1, \xi_2, \dots \rangle \longrightarrow \mathbb{Z}\langle w_1, w_2, \dots \rangle,$$

and its algebraic form is determined by the conditions:

$$\Delta a_* \xi_n = (a_* \otimes a_*)(\Delta \xi_n).$$

For example, $a_* \xi_1 = w_1$, $a_* \xi_2 = w_2 - w_1|w_1$, $a_* \xi_3 = w_3 - 3w_2|w_1 + 2w_1|w_1|w_1$.

Thus using topological results we have obtained that two Hopf algebra structures on the free associative algebra with the comultiplications (4) and (5) are isomorphic over $\mathbb{Z}$.

This result is interesting from the topological point of view, since the elements $(w_n - a_* \xi_n)$ for $n \geq 2$ are obstructions to the desuspension of the homotopy equivalence

$$\Sigma(\Omega\Sigma S^2) \rightarrow \Sigma\left(\bigvee_{n \geq 1} S^{2n}\right).$$

We have the commutative diagram:

$$\begin{array}{ccccc} \Omega\Sigma(\Omega\Sigma S^2) & \xrightarrow{\Omega\Sigma k} & \Omega\Sigma\mathbb{C}P^\infty & & \\ \uparrow & & \uparrow & \searrow j & \\ \Omega\Sigma S^2 & \xrightarrow{k} & \mathbb{C}P^\infty & \xrightarrow{i} & BU \xrightarrow{\det} \mathbb{C}P^\infty \end{array}$$

Here

- i is the inclusion $\mathbb{C}P^\infty = BU(1) \subset BU$,
- j arises from the universal property of $\Omega\Sigma\mathbb{C}P^\infty$ as a free H -space;
- the mapping $\mathbb{C}P^\infty \rightarrow \Omega\Sigma\mathbb{C}P^\infty$ is induced by the identity map $\Sigma\mathbb{C}P^\infty \rightarrow \Sigma\mathbb{C}P^\infty$,
- the mapping $k: \Omega\Sigma S^2 \rightarrow \mathbb{C}P^\infty$ corresponds to the generator of $H^2(\Omega\Sigma S^2) = \mathbb{Z}$.
- $\det$ is the mapping of the classifying spaces $BU \rightarrow BU(1)$ induced by the mappings $\det: U(n) \rightarrow U(1)$, $n = 1, 2, \dots$.

As we have mentioned, $H^*(BU)$ as a Hopf algebra is isomorphic to the algebra of symmetric functions (in the generators – Chern classes) and is self-dual.

$$H^*(BU) \simeq H_*(BU) \simeq \text{Sym}[t_1, t_2, \dots] = \mathbb{Z}[\sigma_1, \sigma_2, \dots].$$

Since $\mathbb{C}P^\infty$ gives rise to the algebra generators in $H_*(BU)$, j_* is an epimorphism on homology, and j^* is a monomorphism on cohomology.

- j_* corresponds to the factorization of the non-commutative polynomial algebra $\mathcal{Z}$ by the commutation relations $Z_i Z_j - Z_j Z_i$ and sends Z_i to σ_i .
- j^* corresponds to the inclusion $\text{Sym}[t_1, t_2, \dots] \subset \text{Qsym}[t_1, t_2, \dots]$ and sends σ_i to M_{ω_i} , where $\omega_i = \underbrace{(1, 1, \dots, 1)}_i$.
- k^* sends u to v_1 and defines an embedding of the polynomial ring $\mathbb{Z}[u]$ into the divided power algebra $\mathbb{Z}[v_n]/I$.
- the composition $\det_* \circ j_*$ maps the algebra $\mathcal{Z}$ to the divided power algebra $H_*(\mathbb{C}P^\infty)$. This corresponds to the mapping $Z_i \rightarrow d_k|_{\mathcal{P}_s}$ of $\mathcal{Z}$ to $\mathcal{D}(\mathcal{P}_s)$.

6 Flag f -vectors

Definition 16. Let P^n be an n -dimensional polytope and $S = \{a_1, \dots, a_k\} \subset \{0, 1, \dots, n-1\}$. A *flag number* $f_S = f_{a_1, \dots, a_k}$ is the number of increasing sequences of faces

$$F^{a_1} \subset F^{a_2} \subset \dots \subset F^{a_k}, \quad \dim F^{a_i} = a_i.$$

It is easy to see that for $S = \{i\}$ the number $f_{\{i\}} = f_i$ is just the number of i -dimensional faces. We have already defined this number above.

The collection $\{f_S\}$ of all the flag numbers is called a *flag f -vector* (or *extended f -vector*) of the polytope P^n . By the definition $f_\emptyset = 1$.

Flag f -vectors have been extensively studied by M. Bayer and L. Billera in [BB], where the generalized Dehn-Sommerville relations are proved:

Theorem. Let P^n be an n -dimensional polytope, and $S \subset \{0, \dots, n-1\}$. If $\{i, k\} \subseteq S \cup \{-1, n\}$ such that $i < k-1$ and $S \cap \{i+1, \dots, k-1\} = \emptyset$, then

$$\sum_{j=i+1}^{k-1} (-1)^{j-i-1} f_{S \cup \{j\}} = (1 - (-1)^{k-i-1}) f_S.$$

Definition 17. For $n \geq 1$ let Ψ^n be the set of subsets $S \subset \{0, 1, \dots, n-2\}$ such that S contains no two consecutive integers.

It is easy to show by induction that the cardinality of Ψ^n is equal to the n -th Fibonacci number c_n ($c_n = c_{n-1} + c_{n-2}$, $c_0 = 1$, $c_1 = 1$).

For any polytope P there exists a *cone* (or a *pyramid*) CP and a *suspension* (or a *bipyramid*) BP . These two operations are defined on combinatorial polytopes and can be extended to linear operators on the ring $\mathcal{P}$. It is natural to set $B\emptyset = 1 = C\emptyset$.

The face lattice of the polytope CP^n is:

$$\{\emptyset\}; \{F_i^0, C\emptyset\}; \{F_i^1, CF_j^0\}; \dots; \{F_i^{n-1}, CF_j^{n-2}\}, \{P^n, CF_i^{n-1}\}; \{CP^n\},$$

where F_i^k are k -dimensional faces of P^n , and

$$\begin{aligned} F^{a_1} \subset F^{a_2} \text{ in } CP &\Leftrightarrow F^{a_1} \subset F^{a_2} \text{ in } P, \\ F^{a_1} \subset CF^{a_2} &\Leftrightarrow F^{a_1} \subset F^{a_2}, \\ CF^{a_1} \subset CF^{a_2} &\Leftrightarrow F^{a_1} \subset F^{a_2}. \end{aligned}$$

The polytope BP^n has the face lattice:

$$\{\emptyset\}, \{C_0\emptyset, C_1\emptyset, F_i^0\}, \{F_i^1, C_0F_j^0, C_1F_k^0\}, \dots, \{F_k^{n-1}, C_0F_j^{n-2}, C_1F_k^{n-2}\}, \{C_0F_i^{n-1}, C_1F_j^{n-1}\}, \{BP^n\}.$$

where C_0 and C_1 are the lower and the upper cones, and

$$\begin{aligned} F^{a_1} \subset F^{a_2} \text{ in } BP &\Leftrightarrow F^{a_1} \subset F^{a_2} \text{ in } P, \\ F^{a_1} \subset C_s F^{a_2} &\Leftrightarrow F^{a_1} \subset F^{a_2}; \\ C_s F^{a_1} \subset C_t F^{a_2} &\Leftrightarrow s = t \text{ and } F^{a_1} \subset F^{a_2}; \end{aligned}$$

Definition 18. For $n \geq 1$ let Ω^n be the set of n -dimensional polytopes that arises when we apply words in B and C that end in C^2 and contain no adjacent B 's to the empty set $\emptyset$.

Each word of length $n+1$ from the set Ω^n either has the form CQ , $Q \in \Omega^{n-1}$, or BCQ , $Q \in \Omega^{n-2}$, so cardinality of the set Ω^n satisfies the Fibonacci relation $|\Omega^n| = |\Omega^{n-1}| + |\Omega^{n-2}|$. Since $|\Omega^1| = |\{C^2\}| = 1$, and $|\Omega^2| = |\{C^3, BC^2\}| = 2$, we see that $|\Omega^n| = c_n = |\Psi^n|$.

M. Bayer and L. Billera proved the following fact:

Theorem. Let $n \geq 1$. Then

1. For all $T \subseteq \{0, 1, \dots, n-1\}$ there is a nontrivial linear relation expressing $f_T(P)$ in terms of $f_S(P)$, $S \in \Psi^n$, which holds for all n -dimensional polytopes (see [BB, Proposition 2.2]).
2. The extended f -vectors of the c_n elements of Ω^n are affinely independent [BB, Proposition 2.3]. Thus the flag f -vectors

$$\{f_S(P^n)\}_{S \in \Psi^n} : P^n - \text{an } n\text{-dimensional polytope}$$

span an $(c_n - 1)$ -dimensional affine hyperplane defined by the equation $f_\emptyset = 1$ (see [BB, Theorem 2.6])

In fact, a little bit more stronger fact which can be easily extracted from the original Bayer-Billera's proof is true.

Let us identify the words in Ω^n with the sets in Ψ^n in such a way that the word $C^{n+1-a_k}BC^{a_k-a_{k-1}-1}B \dots BC^{a_1-1}$ corresponds to the set $\{a_1-3, \dots, a_k-3\}$. Let us set $C < B$ and order the words lexicographically. Consider a matrix K^n of sizes $c_n \times c_n$, $k_{Q,S} = f_S(Q)$, $Q \in \Omega^n$, $S \in \Psi^n$.

Proposition 8.

$$\det(K^n) = 1.$$

Proof. For $n = 1$ the matrix $K^1 = (f_\emptyset(I)) = (1)$.

For $n = 2$

$$K^2 = \begin{pmatrix} 1 & f_0(\Delta^2) \\ 1 & f_0(I^2) \end{pmatrix} = \begin{pmatrix} 1 & 3 \\ 1 & 4 \end{pmatrix}; \quad \det K^2 = 1$$

Let us prove the statement by induction. Consider the matrix K^n :

$$K^n = \begin{pmatrix} K_{11} & K_{12} \\ K_{21} & K_{22} \end{pmatrix}$$

where the block K_{11} corresponds to the words of the form CQ , $Q \in \Omega^{n-1}$ and the sets S , that do not contain $n-2$. Similarly, the block K_{22} corresponds to the words of the form BCQ , $Q \in \Omega^{n-2}$ and the sets containing $n-2$.

Each increasing sequence of faces of the polytope CP^{n-1} without two faces of adjacent dimensions has the form

$$F^{l_1} \subset \dots \subset F^{l_i} \subset CF^{l_{i+1}} \subset \dots \subset CF^{l_k},$$

so for $\{a_1, \dots, a_k\} \in \Psi^n$

$$f_{a_1, \dots, a_k}(CP^{n-1}) = f_{a_1-1, a_2-1, \dots, a_k-1}(P^n) + \dots + f_{a_1, \dots, a_i, a_{i+1}-1, \dots, a_k-1}(P^n) + \dots + f_{a_1, \dots, a_k}(P^n). \quad (6)$$

where $f_{-1, a_2, \dots, a_k} = f_{a_2, \dots, a_k}$.

The sets of the form $\{a_1, \dots, a_l, a_{l+1}-1, \dots, a_k-1\}$ may not belong to Ψ^{n-1} . But we can express them in terms of the sets from Ψ^{n-1} using the generalized Dehn-Sommerville relations. Let $a_k < n-2$. Each relation has the form:

$$f_{S_1, i, k-1, k, S_2} = (-1)^{k-i} \left((1 - (-1)^{k-i-1}) f_{S_1, i, k, S_2} - \sum_{j=i+1}^{k-2} (-1)^{j-i-1} f_{S_1, i, j, k, S_2} \right),$$

where all the sets on the right side are lower than the set on the left side. Thus for $a_k < n-2$

$$f_{a_1, \dots, a_k}(CP^n) = f_{a_1, \dots, a_k}(P^n) + \text{lower summands},$$

so the matrix K_{11} can be represented as $K^{n-1}T$, where T is an upper unitriangular matrix. In particular, $\det K_{11} = \det K^{n-1} \cdot \det T = 1$ by induction.

Lemma 1. *Let P be an $(n-2)$ -dimensional polytope, and $S \in \Psi^n$. Then for the polytopes BCP and CBP*

$$f_S(BCP) = \begin{cases} f_S(CBP), & \text{if } n-2 \notin S, \\ f_S(CBP) + f_{S \setminus \{n-2\}}(P), & \text{if } n-2 \in S, \end{cases}$$

Proof. For the polytope BCP each increasing sequence of faces has one of the forms

$$\begin{aligned} F^{l_1} \subset \dots \subset F^{l_i} \subset CF^{l_{i+1}} \subset \dots \subset CF^{l_j} \subset C_s CF^{l_{j+1}} \subset \dots \subset C_s CF^{l_k} \text{ or} \\ F^{l_1} \subset \dots \subset F^{l_i} \subset C_s F^{l_{i+1}} \subset \dots \subset C_s F^{l_j} \subset C_s CF^{l_{j+1}} \subset \dots \subset C_s CF^{l_k} \text{ or} \\ F^{l_1} \subset \dots \subset F^{l_j} \subset C_s CF^{l_{j+1}} \subset \dots \subset C_s CF^{l_k}, \text{ when } i = j, \end{aligned}$$

where $s = 0$ or 1 .

Similarly, for the polytope CBP each increasing sequence of faces has one of the forms

$$\begin{aligned} F^{l_1} \subset \dots \subset F^{l_i} \subset CF^{l_{i+1}} \subset \dots \subset CF^{l_j} \subset CC_s F^{l_{j+1}} \subset \dots \subset CC_s F^{l_k} \text{ or} \\ F^{l_1} \subset \dots \subset F^{l_i} \subset C_s F^{l_{i+1}} \subset \dots \subset C_s F^{l_j} \subset CC_s F^{l_{j+1}} \subset \dots \subset CC_s F^{l_k} \text{ or} \\ F^{l_1} \subset \dots \subset F^{l_j} \subset CC_s F^{l_{j+1}} \subset \dots \subset CC_s F^{l_k}, \text{ when } i = j, \end{aligned}$$

where $s = 0$ or 1 , except for the additional condition $F^{l_k} \neq P$, that is $l_k \neq n-2$.

Thus for $l_k \neq n-2$, we can exchange CC_s and $C_s C$ to obtain a one-to-one correspondence between the increasing sequences. So $f_S(BCP) = f_S(CBP)$, if $n-2 \notin S$.

If $l_k = n-2$, then we can exchange CC_s and $C_s C$ as before except for the case when the sequence of faces in the polytope BCP has the form

$$F^{a_1} \subset \dots \subset F^{a_{k-1}} \subset P$$

The sequences of this type give exactly $f_{a_1, \dots, a_{k-1}}(P^{n-1})$, so $f_S(BCP) = f_S(CBP) + f_{S \setminus \{n-2\}}(P)$, if $n-2 \in S$. $\square$

Let us consider the polytope BCQ corresponding to one of the lower rows of K^n . If Q starts with C , then there is a row CBQ in the upper part of the matrix. Let us subtract the row CBQ from the row BCQ . Then lemma 1 implies that the resulting row is

$$k'_{BCQ,S} = \begin{cases} 0, & \text{if } n-2 \in S; \\ f_S(Q), & \text{if } n-2 \notin S. \end{cases} \quad (7)$$

If Q starts with B , then there is no row CBQ in the matrix. But since $\det K^{n-1} = 1$ and all flag f -numbers of $(n-1)$ -dimensional polytopes can be expressed in terms of $\{f_S, S \in \Psi^{n-1}\}$, flag f -vector of any $(n-1)$ -dimensional polytope is an integer combination of flag f -vectors of the polytopes from Ω^{n-1} . So flag f -vector of the polytope BQ can be expressed as an integer combination of flag f -vectors of the polytopes $Q' \in \Omega^{n-1}$:

$$f(BQ) = \sum_{Q'} n_{Q'} f(Q')$$

Using the formula (6) we obtain:

$$f(CBQ) = \sum_{Q'} n_{Q'} f(CQ')$$

If we subtract the corresponding integer combination of the rows of the upper part of the matrix from the row BCQ , we obtain the row (7).

Thus we see, that using elementary transformations of rows the matrix K^{n-1} can be transformed to the matrix

$$\begin{pmatrix} K^{n-1}T & K_{12} \\ 0 & K^{n-2} \end{pmatrix}$$

By the inductive assumption $\det K^{n-1} = \det K^{n-2} = 1$, so $\det K^n = 1$. □

Remark. In the proof we follow the original Bayer-Billera's idea, except for the fact that they use the additional matrix of face numbers of the polytopes $Q \in \Psi^n$, and our proof is direct.

Corollary 9. *Flag f -vector of any n -dimensional polytope P^n is an integer combination of flag f -vectors of the polytopes $Q \in \Omega^n$.*

Proof. Indeed, any flag f -number is a linear combination of $f_S, S \in \Psi^n$. Since $\det K^n = 1$, the vector $\{f_S(P), S \in \Psi^n\}$ is an integer combination of the vectors $\{f_S(Q), S \in \Psi^n\}, Q \in \Omega^n$.

This implies that the whole vector $\{f_S(P)\}$ is an integer combination of the vectors $\{f_S(Q)\}, Q \in \Omega^n$ with the same coefficients. □

This fact is important, when we try to describe an image of the generalized f -polynomial (see below) in the ring $\text{Qsym}[t_1, t_2, \dots][\alpha]$.

Generalized f -vectors are also connected with a very interesting construction of the cd -index invented by J.Fine (see [BK, Prop. 2], see also the papers by R. Stanley [St3] and M. Bayer, A. Klapper [BK]).

7 Operators B and C

As it was mentioned below, the operations C and B can be extended to the linear operators on the ring of polytopes.

Then for any n -dimensional polytope P^n we have:

$$\begin{aligned} d_k C P^n &= C d_k P^n + d_{k-1} P^n, \quad 1 \leq k \leq n, \quad d_{n+1} C P^n = 1 + d_n P^n = C \emptyset + d_n P^n; \\ d B P^n &= 2 C P^n, \quad n \geq 1; \quad d B 1 = d I = 2 = 2 C \emptyset, \\ d_k B P^n &= 2 C d_k P^n + d_{k-1} P^n, \quad 2 \leq k \leq n, \quad d_{n+1} B P^n = 2 + d_n P^n = 2 C \emptyset + d_n P^n. \end{aligned}$$

Thus if we add to the ring $\mathcal{P}$ the new element $\emptyset$ of the graduation -1 such that $\emptyset = d1, d\emptyset = 0$, then the operators B and C satisfy the relations:

$$\begin{aligned} dC - Cd &= [d, C] = 1; \quad [d_k, C] = d_{k-1}, \quad k \geq 2; \\ [d, B] &= (2C - B)d; \quad [d_k, B] = (2C - B)d_k + d_{k-1}, \quad k \geq 2. \end{aligned}$$

Otherwise the relations in the ring $\mathcal{P}$ have the form:

$$\begin{aligned} [d, C] &= 1 + \xi_0; \quad [d_{k+1}, C] = d_k + \frac{1}{k!} \frac{\partial^k}{\partial \alpha^k} \Big|_{\alpha=0} \xi_\alpha, \quad k \geq 1; \\ [d, B] &= (2C - B)d + 2\xi_0; \quad [d_{k+1}, B] = (2C - B)d_k + d_{k-1} + \frac{2}{k!} \frac{\partial^k}{\partial \alpha^k} \Big|_{\alpha=0} \xi_\alpha, \quad k \geq 1; \end{aligned}$$

It is convenient to write these relations in terms of the operator $\Phi(t)$. Let us denote $A = 2C - B$. Then

$$\begin{aligned} [\Phi(t), C] &= t\Phi(t) + t\xi_t; \\ [\Phi(t), B] &= (A + t)(\Phi(t) - 1) + 2t\xi_t; \\ \Phi(t)A &= A + t + t\Phi(t); \\ \Phi(t)[B, C] &= [B, C] + tA + t^2; \end{aligned}$$

Proposition 10. *For any $m \geq 0$ the formulas:*

$$\begin{aligned} (Ag)(\alpha, t_1, \dots, t_{m+1}) &= (\alpha + t_{m+1})g(\alpha, 0, \dots, 0) + (t_m + t_{m+1})g(\alpha, t_{m+1}, 0, \dots, 0) + \dots + \\ &\quad + (t_{i-1} + t_i)g(\alpha, t_i, t_{i+1}, \dots, t_{m+1}, 0, \dots, 0) + \dots + t_1(v_{m+1}g)(\alpha, t_1, \dots, t_{m+1}); \\ (Cg)(\alpha, t_1, \dots, t_{m+1}) &= (\alpha + t_1 + \dots + t_{m+1})(v_{m+1}g)(\alpha, t_1, \dots, t_{m+1}) + t_{m+1}g(t_{m+1}, t_1, \dots, t_m) + \\ &\quad + t_m g(t_m, t_1, \dots, t_{m-1}, 0) + \dots + t_i g(t_i, t_1, \dots, t_{i-1}, 0, \dots, 0) + \dots + t_1 g(t_1, 0, \dots, 0) \end{aligned}$$

define the linear mappings $A, C : \text{Qsym}[t_1, \dots, t_m][\alpha] \rightarrow \text{Qsym}[t_1, \dots, t_{m+1}][\alpha]$ such that

$$f_{m+1}(AP^n) = (Af_m)(P^n), \quad \text{and} \quad f_{m+1}(CP^n) = (Cf_m)(P^n)$$

for any $P^n \in \mathcal{P}^n$ and $m \geq n$.

Proof.

$$\begin{aligned} \Phi(t_{m+1})\Phi(t_m) \dots \Phi(t_1)AP &= \Phi(t_{m+1})\Phi(t_m) \dots \Phi(t_2)(A + t_1\Phi(t_1) + t_1)P = \\ &= \Phi(t_{m+1})\Phi(t_m) \dots \Phi(t_3)(A + t_2 + (t_2 + t_1)\Phi(t_2) + t_1\Phi(t_2)\Phi(t_1))P = \\ &= (A + t_{m+1} + (t_{m+1} + t_m)\Phi(t_{m+1}) + (t_m + t_{m-1})\Phi(t_{m+1})\Phi(t_m) + \dots + t_1\Phi(t_{m+1}) \dots \Phi(t_1))P \end{aligned}$$

This gives the formula for $f_{m+1}(AP^n)$, and for $m \geq n$

$$\xi_\alpha \Phi(t_{m+1}) \dots \Phi(t_1)P^n = f_{m+1}(\alpha, t_1, \dots, t_{m+1})(P^n) = (v_{m+1}f_m)(\alpha, t_1, \dots, t_{m+1})(P^n)$$

Similarly we have:

$$\begin{aligned} \Phi(t_{m+1})\Phi(t_m) \dots \Phi(t_1)CP &= \Phi(t_{m+1}) \dots \Phi(t_2)((C + t_1)\Phi(t_1) + t_1\xi_{t_1})P = \\ &= \Phi(t_{m+1})\Phi(t_m) \dots \Phi(t_3)((C + t_2 + t_1)\Phi(t_2)\Phi(t_2) + t_2\xi_{t_2}\Phi(t_1) + t_1\xi_{t_1})P = \\ &= ((C + t_{m+1} + \dots + t_1)\Phi(t_{m+1}) \dots \Phi(t_1)P + t_{m+1}\xi_{t_{m+1}}\Phi(t_m) \dots \Phi(t_1) + \dots + t_1\xi_{t_1})P = \\ &= (C + t_{m+1} + \dots + t_1)\Phi(t_{m+1}) \dots \Phi(t_1)P + t_{m+1}f_m(t_{m+1}, t_1, \dots, t_m)(P) + \\ &\quad + t_m f_m(t_m, t_1, \dots, t_{m-1}, 0)(P) + \dots + t_1 f(t_1, 0, \dots, 0)(P) \end{aligned}$$

Let us prove that for a polynomial $g \in \text{Qsym}[t_1, \dots, t_m][\alpha]$ the polynomials Ag and Cg belong to $\text{Qsym}[t_1, \dots, t_{m+1}][\alpha]$.

$$\begin{aligned} (Ag)(\alpha, t_1, \dots, t_{i-1}, 0, t_{i+1}, \dots, t_{m+1}) &= (\alpha + t_{m+1})g(\alpha, 0, \dots, 0) + \dots + (t_{i+1} + 0)g(\alpha, t_{i+1}, \dots, t_{m+1}, 0, \dots, 0) + \\ &\quad + (0 + t_{i-1})g(\alpha, 0, t_{i+1}, \dots, t_{m+1}, 0, \dots, 0) + \dots + t_1 g(\alpha, t_1, \dots, t_{i-1}, t_{i+1}, t_m) = \\ &= (\alpha + t_{m+1})g(\alpha, 0, \dots, 0) + \dots + (t_{i+1} + t_{i-1})g(\alpha, t_{i+1}, \dots, t_{m+1}, 0, \dots, 0) + \dots + t_{m+1}g(\alpha, t_1, \dots, t_{i-1}, t_{i+1}, \dots, 0) = \\ &= (Ag)(\alpha, t_1, \dots, t_{i-1}, t_{i+1}, \dots, t_{m+1}, 0) \end{aligned}$$

The same situation we have for C :

$$\begin{aligned} (Cg)(\alpha, t_1, \dots, t_{i-1}, 0, t_{i+1}, \dots, t_{m+1}) &= (\alpha + \dots + t_{i-1} + t_{i+1} + \dots + t_{m+1})g(t_{m+1}, t_1, \dots, t_{i-1}, t_{i+1}, \dots, t_{m+1}) + \\ &\quad + \dots + t_{i+1}g(t_{i+1}, t_1, \dots, t_{i-1}, 0, \dots, 0) + 0g(0, t_1, \dots, t_{i-1}, 0, \dots, 0) + t_{i-1}g(t_{i-1}, t_1, \dots, t_{i-2}, 0, \dots, 0) + \\ &\quad + \dots + t_1g(t_1, 0, \dots, 0) = (Cg)(\alpha, t_1, \dots, t_{i-1}, t_{i+1}, \dots, t_{m+1}, 0) \end{aligned}$$

□

The operator B can be defined as $B = 2C - A$. Then $f_{m+1}(BP^n) = (Bf_m)(P^n)$ for $m \geq n$. For the same reasons the operators A, B, C are defined on $\text{Qsym}[t_1, t_2, \dots][\alpha]$:

$$(Ag)(\alpha, t_1, t_2, \dots) = \alpha g(\alpha, 0, \dots) + t_1 g(\alpha, t_1, t_2, \dots) + \sum_{i=2}^{\infty} (t_{i-1} + t_i) g(\alpha, t_i, t_{i+1}, \dots);$$

$$(Cg)(\alpha, t_1, t_2, \dots) = (\alpha + \sigma_1) g(\alpha, t_1, t_2, \dots) + \sum_{i=1}^{\infty} t_i g(t_i, t_1, \dots, t_{i-1}, 0, \dots, 0);$$

$$B = 2C - A,$$

and $(Af)(P) = f(AP)$, $(Bf)(P) = f(BP)$, $(Cf)(P) = f(CP)$.

It is easy to see that

$$\begin{aligned} A(\alpha^m M_{(a_1, \dots, a_k)}) &= \alpha^m (\alpha + 2M_{(1, a_1, \dots, a_k)} + M_{(a_1+1, a_2, \dots, a_k)}), \\ C(\alpha^m M_{(a_1, \dots, a_k)}) &= (\alpha + \sigma_1) M_{(a_1, \dots, a_k)} + M_{(a_1, \dots, a_k, m+1)}. \end{aligned}$$

8 Structure of $\mathcal{D}$

Proposition 11.

$$\Phi(-t)\Phi(t) = 1,$$

that is for any $n \geq 1$

$$d_n - d_1 d_{n-1} + \dots + (-1)^{n-1} d_{n-1} d_1 + (-1)^n d_n = 0 \quad (8)$$

To prove this fact we need the notion of a *face figure*.

Definition 19. Let F be an i -face of the n -dimensional polytope P . A *face figure* P/F is defined as $(F^\diamond)^*$.

The face figure P/F is an $(n - i - 1)$ -dimensional polytope. $(j - i - 1)$ -dimensional faces of P/F are in one-to-one correspondence with j -faces G of P such that $F \subset G \subset P$. Let us denote by G/F the face of P/F corresponding to the face G of P .

Proof.

$$\Phi(-t)\Phi(t) = \sum_{k=0}^{\infty} \left(\sum_{i=0}^k (-1)^i d_i d_{k-i} \right) t^k.$$

Let P^n be an n -dimensional polytope. If $k > n$ then the coefficient of t^k in the series $\Phi(-t)\Phi(t)P^n$ is equal to 0. If $k = 0$, then it is equal to P^n . Let $1 \leq k \leq n$. Then

$$\begin{aligned} (dd_{k-1} - d_2 d_{k-2} + \dots + (-1)^{k-2} d_{k-1} d) P^n &= \\ &= \sum_{F^{n-k} \subset P^n} \left(\sum_{F^{n-k+1} \supset F^{n-k}} 1 - \sum_{F^{n-k+2} \supset F^{n-k}} 1 + \dots + (-1)^{k-2} \sum_{F^{n-1} \supset F^{n-k}} 1 \right) F^{n-k} = \\ &= \sum_{F^{n-k} \subset P^n} (f_0(P^n/F^{n-k}) - f_1(P^n/F^{n-k}) + \dots + (-1)^{k-2} f_{k-2}(P^n/F^{n-k})) F^{n-k} \end{aligned}$$

Since P^n/F^{n-k} is a $(k-1)$ -dimensional polytope, the Euler formula gives the relation

$$f_0(P^n/F^{n-k}) - f_1(P^n/F^{n-k}) + \dots + (-1)^{k-2} f_{k-2}(P^n/F^{n-k}) = (1 + (-1)^k).$$

Thus we obtain

$$(dd_{k-1} - d_2 d_{k-2} + \dots + (-1)^{k-2} d_{k-1} d) P^n = (1 + (-1)^k) d_k P^n \Leftrightarrow \sum_{i=0}^k (-1)^i d_i d_{k-i} P^n = 0$$

□

Theorem 1. *The homomorphism $R : \mathcal{Z} \rightarrow \mathcal{D}$ induces an isomorphism of Hopf algebras:*

$$\mathcal{D} \simeq \mathcal{Z}/J_{\mathcal{D}},$$

where $J_{\mathcal{D}}$ is a two-sided Hopf ideal in the Hopf algebra $\mathcal{Z}$, generated by the relations corresponding to the equality $\Phi(-t)\Phi(t) = 1$.

Proof. Let us prove the following lemma:

Lemma 2. *Let $D \in \mathcal{D}$ be an operator of graduation k . Then for each space $\mathcal{P}^{[n]}$, $n \geq k$ there is a unique representation*

$$D = u(d_2, d_3, \dots, d_k) + dv(d_2, d_3, \dots, d_{k-1}), \quad (9)$$

and there is a unique representation

$$D = u'(d_2, d_3, \dots, d_k) + v'(d_2, \dots, d_{k-1})d, \quad (10)$$

where u, u', v, v' are polynomials.

Proof. For $k = 0$ and 1 it is evident.

Since $2d_2 = d^2$, this is true for $k = 2$. Let $k \geq 3$.

Using the relations (8) we obtain

$$dd_i = (-1)^i d_i d + (d_2 d_{i-1} - d_3 d_{i-2} + \dots + (-1)^{i-1} d_{i-1} d_2) + (1 + (-1)^{i+1}) d_{i+1}$$

So the expressions (9) and (10) exist.

Let $u(d_2, d_3, \dots, d_k) + dv(d_2, \dots, d_{k-1}) = 0$, where $u = \sum_{|\omega|=k} a_{\omega} D_{\omega}$, and $v = \sum_{|\omega|=k-1} b_{\omega} D_{\omega}$.

Then for any n -dimensional polytope P^n , $n \geq k$ we have

$$\xi_1(u + dv)P^n = 0.$$

Since $u + dv$ has degree k , this equality can be written as

$$\sum_{|\omega|=k} a_{\omega} f_{n-k, n-k+j_1, n-k+j_1+j_2, \dots, n-j_l}(P^n) + \sum_{|\omega|=k-1} b_{\omega} f_{n-k, n-k+1, n-k+1+j_1, \dots, n-j_l}(P^n).$$

Using the generalized Dehn-Sommerville equations we obtain

$$f_{n-k, n-k+1, n-k+1+j_1, \dots, n-j_l} = (-1)^{n-k-1} \left(\sum_{j=0}^{n-k-1} (-1)^j f_{j, n-k+1, n-k+1+j_1, \dots, n-j_l} \right) + \\ + (1 + (-1)^{n-k}) f_{n-k+1, n-k+1+j_1, \dots, n-j_l}$$

All the sets $\{j, n-k+1, n-k+1+j_1, \dots, n-j_l\}$ and $\{n-k+1, n-k+1+j_1, \dots, n-j_l\}$ on the right are different for different ω and belong to Ψ^n .

Since the vectors $\{f_S(Q), S \in \Psi^n\}$, $Q \in \Omega^n$ are linearly independent, we obtain that all a_{ω} and b_{ω} are equal to 0, so the representation (9) is unique.

We obtain that the monomials $D_{\omega} = d_{j_1} \dots d_{j_l}$, $|\omega| = k$, $j_i \geq 2$ and $dD_{\omega} = dd_{j_1} \dots d_{j_l}$, $|\omega| = k-1$, $j_i \geq 2$ form a basis of the abelian group $\mathcal{D}^k(\mathcal{P}^{[n]})$.

But each monomial dD_{ω} , $|\omega| = k$, $j_i \geq 2$ can be expressed as an integer combination of the monomials $D_{\omega'}$ and $D_{\omega'}d$. So the monomials D_{ω} and $D_{\omega'}d$ also form a basis. This proves the second part of the lemma. $\square$

Now let us prove the theorem. The mapping $R : \mathcal{Z} \rightarrow \mathcal{D}$ is an epimorphism of Hopf algebras. Let $z \in \mathcal{Z}$ such that $\deg z = k$ and $Rz = 0$.

$$z = \sum_{|\omega|=k} a_{\omega} Z^{\omega}.$$

We know that

$$Z_{k-1}Z_1 = (-1)^{k-1}Z_1Z_{k-1} + \sum_{i=2}^{k-2}(-1)^iZ_{k-i}Z_i + (1 + (-1)^k)Z_k - \sum_{i=0}^k(-1)^iZ_{k-i}Z_i.$$

So

$$z = \sum_{|\omega|=k, j_i \geq 2} a'_\omega Z^\omega + \sum_{|\omega|=k-1, j_i \geq 2} b_\omega Z_1 Z^\omega + z',$$

where $z' \in J_{\mathcal{D}}$. Since $Rz = Rz' = 0$, we obtain

$$\sum_{|\omega|=k, j_i \geq 2} a'_\omega D_{\omega^*} + \sum_{|\omega|=k-1, j_i \geq 2} b_\omega D_{\omega^*} d = 0,$$

Lemma 2 implies that all the coefficients a_ω and b_ω are equal to 0, so $z = z' \in J_{\mathcal{D}}$. This proves that $\mathcal{D} \simeq \mathcal{Z}/J_{\mathcal{D}}$. $\square$

It is evident that $D_\omega|_{\mathcal{P}^{[n]}} = 0$ for $|\omega| > n$. The following corollary says that the other relations between operators in $\mathcal{D}$ on the abelian group $\mathcal{P}^{[n]}$ are the same as on the whole ring $\mathcal{P}$.

Corollary 12. *We have*

$$\mathcal{D}(\mathcal{P}^{[n]}) = \mathcal{D}/J_n$$

where the ideal J_n is generated by all the operators D_ω , $|\omega| > n$.

Proof. As in the proof of Theorem 1 using Lemma 2 it is easy to show that $\mathcal{D}(\mathcal{P}^{[n]}) \simeq \mathcal{Z}/J_{\mathcal{D}(\mathcal{P}^{[n]})}$, where the ideal $J_{\mathcal{D}(\mathcal{P}^{[n]})}$ is generated by the relations $\Phi(t)\Phi(-t) = 1$ and $Z_\omega = 0$ for $|\omega| > n$. $\square$

Corollary 13. *The operators $d_2, d_3, d_4, \dots$ are algebraically independent.*

Corollary 14. *Rank of the n -th graded component of the ring $\mathcal{D}$ is equal to the $(n-1)$ -th Fibonacci number c_{n-1} .*

Proof. At first, let us calculate rank r_n of the n -th graded component of the ring $\mathbb{Z}\langle d_2, d_3, d_4, \dots \rangle$.

$$r_0 = 1, r_1 = 0, r_2 = 1, r_3 = 1, r_4 = 2, r_5 = 3, \dots$$

It is easy to see that there is a recursive relation

$$r_{n+1} = r_{n-1} + r_{n-2} + \dots + r_2 + 1, n \geq 3$$

Then $r_{n+1} = r_{n-1} + r_n$. Since $r_2 = r_3 = 1$ we obtain $r_n = c_{n-2}$, $n \geq 2$. At last, rank of the n -th graded component of the ring $\mathcal{D}$ is equal to $r_{n-1} + r_n = c_{n-3} + c_{n-2} = c_{n-1}$ for $n \geq 3$. It is easy to see that for $n = 1$ and 2 it is also true. $\square$

We claim that the antipode in $\mathcal{D}$ is $\chi : d_k \rightarrow (-1)^k d_k$. Indeed, the relations $\Phi(-t)\Phi(t) = 1$ exactly mean that χ is an antipode. Therefore we obtain the corollary:

Corollary 15. *The Hopf algebra $\mathcal{D}$ is a universal Hopf algebra in the category of Leibnitz-Hopf algebras with an antipode satisfying the formula $\chi(H_i) = (-1)^i H_i$.*

Proof. In fact, let us consider the equation (2) with $\chi(Z_i) = (-1)^i Z_i$:

$$(-1)^{n+1}Z_{n+1} + (-1)^n Z_n Z_1 + \dots + (-1)Z_1 Z_n + Z_{n+1} = 0.$$

The collection of these equalities is equivalent to the equation

$$\Phi(-t)\Phi(t) = 1;$$

$\square$

Let us remind that on the ring of simple polytopes $d_k|_{\mathcal{P}_s} = \frac{d^k}{k!}|_{\mathcal{P}_s}$, so the ring $\mathcal{D}(\mathcal{P}_s)$ is isomorphic to the divided power ring $d_k d_l = \binom{k+l}{k} d_{k+l}$.

Proposition 16.

$$\mathcal{D} \otimes \mathbb{Q} = \mathbb{Q}\langle d_1, d_3, d_5, \dots \rangle,$$

The inclusion $\mathcal{D} \subset \mathcal{D} \otimes \mathbb{Q}$ is an embedding, and the operators d_{2k} are expressed in terms of the operators $d_1, d_3, \dots, d_{2k-1}$ by the formulas

$$d_{2k} = \sum_{i=1}^k (-1)^{i-1} \frac{\binom{2i-2}{i-1}}{i 2^{2i-1}} \left(\sum_{j_1+j_2+\dots+j_{2i}=i+k} d_{2j_1-1} \dots d_{2j_{2i}-1} \right). \quad (11)$$

Proof. Let us denote

$$a(t) = \sum_{k \geq 0} d_{2k} t^{2k}, \quad b(t) = \sum_{k \geq 0} d_{2k+1} t^{2k+1}.$$

Then

$$\begin{aligned} \Phi(t) &= a(t) + b(t) & \Phi(-t) &= a(t) - b(t); \\ a(t)b(t) &= \frac{\Phi(t) + \Phi(-t)}{2} \frac{\Phi(t) - \Phi(-t)}{2} = \frac{\Phi(t)^2 - \Phi(-t)^2}{4} = b(t)a(t). \end{aligned}$$

Then the relation $\Phi(-t)\Phi(t) = 1$ is equivalent to the relations

$$\begin{aligned} a(t)^2 - b(t)^2 &= 1; \\ a(t)b(t) &= b(t)a(t). \end{aligned}$$

Therefore $a(t) = \sqrt{1 + b(t)^2}$ and the formula (11) is true. Consequently all the operators d_{2k} are expressed as polynomials in $d_1, d_3, \dots$ with rational coefficients. For example,

$$d_2 = \frac{d^2}{2}, \quad d_4 = \frac{dd_3 + d_3d}{2} - \frac{d^4}{8}.$$

This means that the algebra $\mathcal{D} \otimes \mathbb{Q}$ is generated by $d_1, d_3, d_5, \dots$. On the other hand, let us calculate the number of the monomials $d_{2j_1-1} \dots d_{2j_k-1}$ with $(2j_1-1) + \dots + (2j_k-1) = n$. Denote this number by l_n . Then

$$l_0 = 1, \quad l_1 = 1, \quad l_2 = 1, \quad l_3 = 2, \quad l_4 = 3, \dots,$$

and there is a recursive formula $l_{n+1} = l_n + l_{n-2} + l_{n-4} + \dots$ for $n \geq 2$. Hence $l_{n+1} = l_n + l_{n-1}$ for $n \geq 2$, and $l_n = c_{n-1}$ for $n \geq 1$.

We see that the number of monomials of degree n is equal to dimension of the n -th graded component according to Corollary 14. This implies that they are linearly independent over the rationals. Therefore $d_1, d_3, d_5, \dots$ are algebraically independent. $\square$

Definition 20. Let us define operators s_k by the formula

$$s(t) = s_1 t + s_2 t^2 + s_3 t^3 + \dots = \log \Phi(t).$$

Then $s_1 = d_1$, $s_2 = 0$, $s_3 = d_3 - \frac{d^3}{6}$, and so on.

The relation $\Phi(-t)\Phi(t) = 1$ turns into the relation $s(-t) + s(t) = 0$. Thus $s_{2k} = 0$ for all k .

Also we have $\Delta s(t) = 1 \otimes s(t) + s(t) \otimes 1$, so each operator s_{2k-1} is a derivation.

Proposition 17. *There is an isomorphism of Hopf algebras*

$$\mathcal{D} \otimes \mathbb{Q} = \mathbb{Q}\langle s_1, s_3, s_5, \dots \rangle,$$

where $\mathbb{Q}\langle s_1, s_3, s_5, \dots \rangle$ is a free Lie-Hopf algebra in the generators of odd degree, the comultiplication $\Delta s_{2k-1} = 1 \otimes s_{2k-1} + s_{2k-1} \otimes 1$, and the antipode $\chi(s_{2k-1}) = -s_{2k-1}$

9 Generalized f -polynomial

The mapping $\Phi : \mathcal{P} \rightarrow \mathcal{P}[t]$ is a ring homomorphism. If we set $\Phi(t_2)t_1 = t_1$, then the operator $\Phi(t_2)$ defines a ring homomorphism $\mathcal{P} \rightarrow \mathcal{P}[t_1, t_2] : P \rightarrow \Phi(t_2)\Phi(t_1)P$,

Thus for each $n \geq 0$ we can define the ring homomorphism $\Phi_n : \mathcal{P} \rightarrow \mathcal{P}[[t_1, \dots, t_n]]$ as a composition:

$$\Phi_n(t_1, \dots, t_n)P = \Phi(t_n) \dots \Phi(t_1)P.$$

Definition 21. Let us define the mapping $\varphi : \mathcal{P} \rightarrow \mathcal{D}^*[\alpha]$:

$$P \rightarrow \varphi_{\alpha, P}, \quad \varphi_{\alpha, P}(D) = \xi_{\alpha}(DP),$$

where $\mathcal{D}^*$ is a graded dual Hopf algebra, and $P \in \mathcal{P}$, $D \in \mathcal{D}$.

It is easy to see that φ is a ring homomorphism. This follows from the equality

$$D_{\omega}(P \times Q) = \mu \Delta D_{\omega}(P \otimes Q),$$

which holds for all compositions ω and all polytopes P, Q . Here μ is the multiplication in the ring of polytopes.

Definition 22. Let P^n be an n -dimensional polytope. Let us define a quasi-symmetric function

$$f_m(\alpha, t_1, \dots, t_m)(P^n) = \xi_{\alpha} \Phi(t_m) \dots \Phi(t_1)P^n = \alpha^n + \sum_{k=1}^{\min\{m, n\}} \sum_{0 \leq a_1 < \dots < a_k \leq n-1} f_{a_1, \dots, a_k} \alpha^{a_1} M_{(n-a_k, \dots, a_2-a_1)}.$$

For $n = 1$ we obtain $f_1(\alpha, t_1)(P^n) = \alpha^n + \sum_{i=1}^n f_i \alpha^i t_1^{n-i}$ is a homogeneous f -polynomial in two variables ([Buch]). So the polynomial f_n is a generalization of the f -polynomial.

Consider the increasing sequence of rings

$$\mathbb{Z}[\alpha] \subset \mathbb{Z}[\alpha, t_1] \subset \mathbb{Z}[\alpha, t_1, t_2] \subset \dots$$

with the restriction maps $u_m : \mathbb{Z}[\alpha, t_1, \dots, t_{m+1}] \rightarrow \mathbb{Z}[\alpha, t_1, \dots, t_m]$

$$(u_m g)(\alpha, t_1, \dots, t_m) = g(\alpha, t_1, \dots, t_m, 0).$$

Since $f_{m+1}(\alpha, t_1, \dots, t_m, 0)(P^n) = f_m(\alpha, t_1, \dots, t_m)(P^n)$, we obtain the ring homomorphism

$$f : \mathcal{P} \rightarrow \text{Qsym}[t_1, t_2, \dots][\alpha] \subset \varprojlim_m \mathbb{Z}[\alpha, t_1, \dots, t_m] :$$

$$f(\alpha, t_1, t_2, \dots)(P^n) = \alpha^n + \sum_{k=1}^n \sum_{0 \leq a_1 < \dots < a_k \leq n-1} f_{a_1, \dots, a_k} \alpha^{a_1} M_{(n-a_k, \dots, a_2-a_1)}.$$

It follows from the formula, that the restriction

$$f(\alpha, t_1, \dots, t_m, t_{m+1}, \dots) \rightarrow f(\alpha, t_1, \dots, t_m, 0, 0, \dots) = f_m(\alpha, t_1, \dots, t_m)$$

is injective on the space of all n -dimensional polytopes, $n \leq m$.

Theorem 2. The image of the space $\mathcal{P}^n$ generated by all n -dimensional polytopes in the ring $\text{Qsym}[t_1, \dots, t_m][\alpha]$, $m \geq n$ under the mapping f_m consists of all the homogeneous polynomials of degree n satisfying the equations

1.

$$\begin{aligned} f(\alpha, t_1, -t_1, t_3, \dots, t_m) &= f(\alpha, 0, 0, t_3, \dots, t_m); \\ f(\alpha, t_1, t_2, -t_2, t_4, \dots, t_m) &= f(\alpha, t_1, 0, 0, t_4, \dots, t_m); \\ &\dots \\ f(\alpha, t_1, \dots, t_{m-2}, t_{m-1}, -t_{m-1}) &= f(\alpha, t_1, \dots, t_{m-2}, 0, 0); \end{aligned}$$

2.

$$f(-\alpha, t_1, \dots, t_{m-1}, \alpha) = f(\alpha, t_1, \dots, t_{m-1}, 0);$$

These equations are equivalent to the Bayer-Stillera (generalized Dehn-Sommerville) relations.

Proof. $\Phi(-t)\Phi(t) = 1 = \Phi(0)\Phi(0)$, therefore

$$\begin{aligned} \xi_\alpha \Phi(t_m) \dots \Phi(t_3) \Phi(-t_1) \Phi(t_1) P^n &= \xi_\alpha \Phi(t_m) \dots \Phi(t_3) \Phi(0) \Phi(0) P^n; \\ \xi_\alpha \Phi(t_m) \dots \Phi(t_4) \Phi(-t_2) \Phi(t_2) \Phi(t_1) P^n &= \xi_\alpha \Phi(t_m) \dots \Phi(t_4) \Phi(0) \Phi(0) \Phi(t_1) P^n; \\ &\dots \\ \xi_\alpha \Phi(-t_{m-1}) \Phi(t_{m-1}) \Phi(t_{m-2}) \dots \Phi(t_1) P^n &= \xi_\alpha \Phi(0) \Phi(0) \Phi(t_{m-2}) \dots \Phi(t_1) P^n. \end{aligned}$$

On the other hand, Proposition 3 gives the last relation

$$\xi_{-\alpha} \Phi(\alpha) \Phi(t_{m-1}) \dots \Phi(t_1) P^n = \xi_\alpha \Phi(0) \Phi(t_{m-1}) \dots \Phi(t_1) P^n.$$

Now let us proof the opposite inclusion, that is if the homogeneous polynomial g of degree n satisfies the conditions of the theorem, then $g = f_m(p^n)$ for some $p^n \in \mathcal{P}^n$.

Lemma 3. *Let P^n be an n -dimensional polytope. Then the equation*

$$f_m(\alpha, t_1, \dots, t_q, -t_q, \dots, t_m) = f_m(\alpha, t_1, \dots, 0, 0, \dots, t_m)$$

is equivalent to the generalized Dehn-Sommerville relations

$$\sum_{j=a_t}^{a_{t+1}-1} (-1)^{j-a_t-1} f_{a_1, \dots, a_t, j, a_{t+1}, \dots, a_k} = (1 + (-1)^{a_{t+1}-a_t}) f_{a_1, \dots, a_t, a_{t+1}, \dots, a_k}$$

for $1 \leq k \leq \min(m-1, n-1)$, $k+1-q \leq t \leq m-q$.

The equation

$$f_m(-\alpha, t_1, \dots, t_{m-1}, \alpha) = f_m(\alpha, t_1, \dots, t_{m-1}, 0)$$

is equivalent to the generalized Dehn-Sommerville relations

$$\sum_{j=0}^{a_1-1} f_{j, a_1, \dots, a_k} = (1 + (-1)^{a_1-1}) f_{a_1, \dots, a_k}$$

for $0 \leq k \leq \min(m-1, n-1)$.

Proof.

$$\begin{aligned} f_m(\alpha, t_1, \dots, t_q, -t_q, \dots, t_m) &= \alpha^n + \sum_{k=1}^{\min(m, n)} \sum_{0 \leq a_1 < \dots < a_k \leq n-1} f_{a_1, \dots, a_k} \alpha^{a_1} \left(\sum_{1 \leq l_1 < \dots < l_k \leq m} t_{l_1}^{n-a_k} \dots t_{l_k}^{a_2-a_1} \right) = \\ &= \alpha^n + \sum_{k=1}^{\min(m-2, n)} \sum_{0 \leq a_1 < \dots < a_k \leq n-1} f_{a_1, \dots, a_k} \alpha^{a_1} \left(\sum_{1 \leq l_1 < \dots < l_k \leq m, l_j \neq q, q+1} t_{l_1}^{n-a_k} \dots t_{l_k}^{a_2-a_1} \right) + \\ &+ \sum_{k=1}^{\min(m-1, n)} \sum_{0 \leq a_1 < \dots < a_k \leq n-1} f_{a_1, \dots, a_k} \alpha^{a_1} \left(\sum_{1 \leq l_1 < \dots < l_j = q < \dots < l_k \leq m, l_{j+1} \neq q+1} t_{l_1}^{n-a_k} \dots t_q^{a_{k+2-j}-a_{k+1-j}} \dots t_{l_k}^{a_2-a_1} \right) + \\ &+ \sum_{k=1}^{\min(m-1, n)} \sum_{0 \leq a_1 < \dots < a_k \leq n-1} f_{a_1, \dots, a_k} \alpha^{a_1} \left(\sum_{1 \leq l_1 < \dots < l_{j+1} = q+1 < \dots < l_k \leq m, l_j \neq q} t_{l_1}^{n-a_k} \dots (-t_q)^{a_{k+1-j}-a_{k-j}} \dots t_{l_k}^{a_2-a_1} \right) + \\ &\sum_{k=1}^{\min(m, n)} \sum_{0 \leq a_1 < \dots < a_k \leq n-1} f_{a_1, \dots, a_k} \alpha^{a_1} \left(\sum_{1 \leq l_1 < \dots < l_j = q < l_{j+1} = q+1 < \dots < l_k \leq m} t_{l_1}^{n-a_k} \dots (-1)^{a_{k+1-j}-a_{k-j}} t_q^{a_{k+2-j}-a_{k-j}} \dots t_{l_k}^{a_2-a_1} \right) \end{aligned}$$

The first summand is exactly $f(\alpha, t_1, \dots, 0, 0, \dots, t_m)$. Therefore all the coefficients of the polynomial consisting of the last three summands should be equal to 0.

Consider the monomial $\alpha^{a_1} t_{l_1}^{n-a_k} \dots t_q^{a_{t+1}-a_t} \dots t_{l_k}^{a_2-a_1}$. Here $q = l_{k+1}-t$. The existence of a monomial of this form in the sum is equivalent to the conditions

$$k \leq \min(m-1, n-1), \quad k+1-t \leq q, \quad t-1 \leq m-q-1.$$

The coefficient of the monomial should be equal to 0. This is equivalent to the relation:

$$(1 + (-1)^{a_{t+1}-a_t}) f_{a_1, \dots, a_t, a_{t+1}, \dots, a_k} + \sum_{j=a_t+1}^{a_{t+1}-1} (-1)^{j-a_t} f_{a_1, \dots, a_t, j, a_{t+1}, a_k} = 0$$

Now let us consider the remaining relation $f(-\alpha, t_1, \dots, t_{m-1}, \alpha) = f(\alpha, t_1, \dots, t_{m-1}, 0)$:

$$\begin{aligned} &= (-\alpha)^n + \sum_{k=1}^{\min(m-1, n)} \sum_{0 \leq a_1 < \dots < a_k \leq n-1} f_{a_1, \dots, a_k} (-\alpha)^{a_1} \left(\sum_{1 \leq l_1 < \dots < l_k \leq m-1} t_{l_1}^{n-a_k} \dots t_{l_k}^{a_2-a_1} \right) + \\ &+ \sum_{k=1}^{\min(m, n)} \sum_{0 \leq a_1 < \dots < a_k \leq n-1} f_{a_1, \dots, a_k} (-\alpha)^{a_1} \left(\sum_{1 \leq l_1 < \dots < l_k = m} t_{l_1}^{n-a_k} \dots t_{l_{k-1}}^{a_3-a_2} \alpha^{a_2-a_1} \right) = \\ &\alpha^n + \sum_{k=1}^{\min(m-1, n)} \sum_{0 \leq a_1 < \dots < a_k \leq n-1} f_{a_1, \dots, a_k} \alpha^{a_1} \left(\sum_{1 \leq l_1 < \dots < l_k \leq m-1} t_{l_1}^{n-a_k} \dots t_{l_k}^{a_2-a_1} \right) \end{aligned}$$

This is equivalent to the relations

$$(-1)^{a_1} f_{a_1, \dots, a_k} + \sum_{j=0}^{a_1-1} (-1)^j f_{j, a_1, \dots, a_k} = f_{a_1, \dots, a_k}$$

for $0 \leq k \leq \min(m-1, n-1)$. If $k=0$, then the corresponding relation is exactly the Euler formula

$$(-1)^n + (-1)^{n-1} f_{n-1} + \dots + f_2 - f_1 + f_0 = 1.$$

□

Corollary 18. For $m \geq n$ relations 1. and 2. of the theorem are equivalent to the generalized Dehn-Sommerville relations for the polytope P^n : For $S \subset \{0, \dots, n-1\}$, and $\{i, k\} \subseteq S \cup \{-1, n\}$ such that $i < k-1$ and $S \cap \{i+1, \dots, k-1\} = \emptyset$:

$$\sum_{j=i+1}^{k-1} (-1)^{j-i-1} f_{S \cup \{j\}} = (1 - (-1)^{k-i-1}) f_S.$$

Proof. We see that relations 1. and 2. follow from the generalized Dehn-Sommerville relations.

On the other hand, if $i = -1$ then the corresponding relation follows from equation 2.

If $S = \{a_1, \dots, a_t, a_{t+1}, \dots, a_s\}$, $i = a_t \geq 0$, and $k = a_{t+1}$; or $S = \{a_1, \dots, a_t\}$, $i = a_t$, and $k = n$, then we can take q such that $s+1-t \leq q \leq m-t$. □

Remark. For n -dimensional polytopes for different $m \geq n$ not all the equations are independent. In fact, for $i = -1$ relation (6) follows from equation 2. and all the equations of type 1. do not contain the case $i = -1$.

On the other hand, let $S = \{a_1, \dots, a_t, a_{t+1}, \dots, a_s\}$, $i = a_t \geq 0$, and $k = a_{t+1}$; or $S = \{a_1, \dots, a_t\}$, $i = a_t$, and $k = n$. Lemma 3 implies that relation (6) follows from the equation with $s+1-t \leq q \leq m-t$. Let us denote $a = s+t-1$, $b = m-t$. There are two conditions for s, t , namely

$$1 \leq s \leq n-1 \text{ and } 1 \leq t \leq s.$$

Let us rewrite these conditions in terms of a and b : $t = m - b$; $s = a + t - 1 = a - b + m - 1$, so

$$m - n \leq b - a \leq m - 2; \quad b \leq m - 1; \quad 1 \leq a$$

This gives us a triangle on the plane (a, b) . Each point (a, b) of this triangle corresponds to the conditions $a \leq q \leq b$, that is there should be the equation for q in the segment $[a, b]$. We can imagine that this segment is the segment $[(a, a), (a, b)]$ on the plane.

Thus for $m = n$ all the equations for $t_1, \dots, t_{n-1}$ are necessary, for $m = n + 1$ it is enough to take the equations for $q = 2, 4, 6, \dots, 2[\frac{n}{2}]$. If $m \geq 2n - 2$ one equation

$$f(\alpha, t_1, \dots, t_{n-1}, -t_{n-1}, t_{n+1}, \dots, t_m) = f(\alpha, t_1, \dots, 0, 0, t_{n+1}, \dots, t_m)$$

gives all the relations of type 1.

Now let us finish the proof of the theorem. If the homogeneous polynomial $g \in \text{Qsym}[t_1, \dots, t_m][\alpha]$ of degree n satisfies all the relations of the theorem, then its coefficients satisfy the generalized Dehn-Sommerville relations. Therefore all the coefficients are linear combinations of the coefficients $g_{a_1, \dots, a_k}$, where $S = \{a_1, \dots, a_k\} \in \Psi^n$. As we know, the vectors $\{f_S(Q), S \in \Psi^n\}$, $Q \in \Omega^n$ form a basis of the abelian group of all the vectors $\{f_S, S \in \Psi^n, f_S \in \mathbb{Z}\} = \mathbb{Z}^{c_n}$. So the vector $\{g_S, S \in \Psi^n\}$ is an integer combination of the vectors $\{f_S(Q), S \in \Psi^n\}$, $Q \in \Omega^n$. This implies that the polynomial g is an integer combination of the polynomials $f_m(Q)$, $Q \in \Omega^n$ with the same coefficients. $\square$

Let us remind that rank of the space $f_m(\mathcal{P}^n)$, $m \geq n$ is equal to c_n .

Proposition 19. *Let $m \geq 2$. Then $f_m(\alpha, t_1, \dots, t_m)(P^n) = f_1(\alpha, t_1 + \dots + t_m)(P^n)$ if and only if P^n is a simple polytope. Here $f_1(\alpha, t)$ is a usual homogeneous f -polynomial in two variables.*

Proof. On the ring of simple polytopes $d_k|_{\mathcal{P}_s} = \frac{d^k}{k!}|_{\mathcal{P}_s}$, so $\Phi(t) = e^{td}$.

Then $\Phi(t_m)\Phi(t_{m-1})\dots\Phi(t_1) = \Phi(t_1 + \dots + t_m)$, therefore $f_m(\alpha, t_1, \dots, t_m)(P^n) = f_1(\alpha, t_1 + \dots + t_m)(P^n)$.

On the other hand, let $f_m(\alpha, t_1, \dots, t_m)(P^n) = f_1(\alpha, t_1 + \dots + t_m)(P^n)$.

Then $f_2(\alpha, t_1, t_2)(P^n) = f_1(\alpha, t_1 + t_2)(P^n)$. So

$$\alpha^n + \sum_{i=0}^{n-1} f_i \alpha^i (t_1^{n-i} + t_2^{n-i}) + \sum_{0 \leq i < j \leq n-1} f_{ij} \alpha^i t_1^{n-j} t_2^{j-i} = \alpha^n + \sum_{i=0}^{n-1} f_i \alpha^i (t_1 + t_2)^{n-i}$$

In particular, $f_{01} = n f_0$, since the coefficients of the monomial $t_1^{n-1} t_2$ on the left and on the right are equal. Hence $2f_1 = f_{01} = n f_0$. This implies that the polytope P^n is simple. $\square$

Remark. Letting m tend to infinity we obtain that $f(\alpha, t_1, t_2, \dots)(P^n) = f_1(\alpha, t_1 + t_2 + \dots)(P^n)$ if and only if P^n is simple.

In the case of simple polytopes the equations of the first type are trivial, but the equation of the second type has the form:

$$f_1(-\alpha, t_1 + \dots + t_{m-1} + \alpha) = f_1(\alpha, t_1 + \dots + t_{m-1})$$

If we denote $t = t_1 + \dots + t_{m-1}$, then $f_1(-\alpha, \alpha + t) = f_1(\alpha, t)$. This equation is equivalent to the Dehn-Sommerville relations (after the change of variables $h(\alpha, t) = f_1(\alpha - t, t)$ it is equivalent to the fact that $h(\alpha, t) = h(t, \alpha)$)

10 Characterization of f

In this part we find the condition that uniquely determines the generalized f -polynomial.

At first let us find the relation between $f(P^n)$ and $f(d_k P^n)$.

Proposition 20. *For any polytope $P^n \in \mathcal{P}$ we have*

$$f(\alpha, t_1, t_2, \dots)(d_k P^n) = \frac{1}{k!} \frac{\partial^k}{\partial t^k} \Big|_{t=0} f(\alpha, t, t_1, t_2, \dots)(P^n)$$

Proof. Indeed,

$$\begin{aligned} \frac{1}{k!} \frac{\partial^k}{\partial t^k} \Big|_{t=0} f(\alpha, t, t_1, t_2, \dots)(P^n) &= \frac{1}{k!} \frac{\partial^k}{\partial t^k} \Big|_{t=0} \lim_{r \rightarrow \infty} \xi_\alpha \Phi(t_r) \dots \Phi(t_1) \Phi(t) P^n = \\ \lim_{r \rightarrow \infty} \frac{1}{k!} \frac{\partial^k}{\partial t^k} \Big|_{t=0} \xi_\alpha \Phi(t_r) \dots \Phi(t_1) \Phi(t) P^n &= \lim_{r \rightarrow \infty} \xi_\alpha \Phi(t_r) \dots \Phi(t_1) d_k P^n = f(\alpha, t_1, t_2, \dots)(d_k P^n) \end{aligned}$$

□

Corollary 21.

$$\begin{aligned} f(\alpha, t, t_1, t_2, \dots)(P^n) &= f(\alpha, t_1, t_2, \dots)(P^n) + f(\alpha, t_1, t_2, \dots)(dP^n)t + \\ &\quad + f(\alpha, t_1, t_2, \dots)(d_2 P^n)t^2 + \dots + f(\alpha, t_1, t_2, \dots)(d_n P^n)t^n \\ f_{m+1}(\alpha, t, t_1, \dots, t_m)(P^n) &= f_m(\alpha, t_1, \dots, t_m)(P^n) + f_m(\alpha, t_1, \dots, t_m)(dP^n)t + \\ &\quad + f_m(\alpha, t_1, \dots, t_m)(d_2 P^n)t^2 + \dots + f_m(\alpha, t_1, \dots, t_m)(d_n P^n)t^n, \quad m \geq 0; \end{aligned}$$

Proof. The first equality follows from Proposition 20.

In fact, both equalities can be proved directly:

$$\begin{aligned} f_{m+1}(\alpha, t, t_1, \dots, t_m)(P^n) &= \xi_\alpha \Phi(t_m) \dots \Phi(t_1) \Phi(t) P^n = \sum_{k=0}^n (\xi_\alpha \Phi(t_m) \dots \Phi(t_1) d_k P^n) t^k = \\ &= \sum_{k=0}^n f_m(\alpha, t_1, \dots, t_m)(d_k P^n) t^k. \end{aligned}$$

Letting m tend to infinity we obtain the first equality. □

The first equality of Corollary 21 is equivalent to the condition $f(\alpha, t_1, t_2, \dots)(\Phi(t)P^n) = f(\alpha, t, t_1, t_2, \dots)(P^n)$. Thus we see that the following diagram commutes

$$\begin{array}{ccc} \mathcal{P} & \xrightarrow{f} & \text{Qsym}[t_1, t_2, \dots][\alpha] \\ \Phi \downarrow & & T \downarrow \\ \mathcal{P}[t] & \xrightarrow{f} & \text{Qsym}[t, t_1, t_2, \dots][\alpha] \end{array}$$

where $f(t) = t$, and $T : \text{Qsym}[t_1, t_2, \dots][\alpha] \rightarrow \text{Qsym}[t, t_1, t_2, \dots][\alpha]$ is a ring homomorphism:

$$Tg(\alpha, t_1, t_2, \dots) = g(\alpha, t, t_1, t_2, \dots), \quad g \in \text{Qsym}[t_1, t_2, \dots][\alpha].$$

Remark. Note that $f_m(\alpha, t_1, \dots, t_m)(\Phi(t)P^n) = f_{m+1}(\alpha, t, t_1, \dots, t_m)(P^n)$ for all $m \geq 0$.

Consider the ring homomorphism $T_{m+1} : \text{Qsym}[t_1, \dots, t_m][\alpha] \rightarrow \text{Qsym}[t, t_1, \dots, t_m][\alpha]$ defined as

$$T_{m+1}(\alpha) = \alpha, \quad T_{m+1}M_\omega(t_1, \dots, t_m) = M_\omega(t, t_1, \dots, t_m)$$

Then the corresponding diagram commutes only for $m \geq n$.

Theorem 3. Let $\psi : \mathcal{P} \rightarrow \text{Qsym}[t_1, t_2, \dots][\alpha]$ be a linear map such that

1. $\psi(\alpha, 0, 0, \dots)(P^n) = \alpha^n$;
2. The following diagram commutes:

$$\begin{array}{ccc} \mathcal{P} & \xrightarrow{\psi} & \text{Qsym}[t_1, t_2, \dots][\alpha] \\ \Phi \downarrow & & T \downarrow \\ \mathcal{P}[t] & \xrightarrow{\psi} & \text{Qsym}[t, t_1, t_2, \dots][\alpha] \end{array}$$

Then $\psi = f$.

Proof. The first condition implies that $\psi(\alpha, 0, 0, \dots)(p) = \xi_\alpha p$.

Let $\psi(P^n) = \alpha^n + \sum_{\omega} \psi_{\omega}(\alpha) M_{\omega}$, and let $\omega = (j_1, \dots, j_k)$.

Since $\psi(\alpha, t, t_1, t_2, \dots)(P^n) = \psi(\alpha, t_1, t_2, \dots)(\Phi(t)P^n)$, we have

$$\psi(\alpha, y_1, y_2, \dots, y_k, t_1, t_2, \dots)(P^n) = \psi(\alpha, t_1, t_2, \dots)(\Phi(y_k) \dots \Phi(y_1)P^n)$$

Therefore

$$\psi(\alpha, y_1, \dots, y_k, 0, 0, \dots) = \psi(\alpha, 0, 0, \dots)(\Phi(y_k) \dots \Phi(y_1)P^n) = \xi_\alpha \Phi(y_k) \dots \Phi(y_1)P^n = f(\alpha, y_1, \dots, y_k, 0, 0, \dots)$$

Hence $\psi_{\omega}(\alpha) = f_{\omega} \alpha^{n-|\omega|}$ for all ω , so $\psi = f$. $\square$

Remark. Let us mention that the mapping T is an isomorphism $\text{Qsym}[t_1, t_2, \dots][\alpha] \rightarrow \text{Qsym}[t, t_1, t_2, \dots][\alpha]$, while Φ is an injection and its image is described by the condition: $p(t) \in \Phi(\mathcal{P})$ is and only if $\Phi(-t)p(t) \in \mathcal{P}$.

On the ring of simple polytopes we have $f(\alpha, t_1, t_2, \dots)(p) = f_1(\alpha, t_1 + t_2 + \dots)(p) = f_1(\alpha, \sigma_1)(p)$, so the image of $\mathcal{P}$ belongs to $\mathbb{Z}[\alpha, \sigma_1]$. On the other hand, $\Phi(t) = e^{td}$, so we have the condition $f_1(\alpha, t_1)(e^{td}p) = f(\alpha, t + t_1)(p)$.

Proposition 22. Let $\psi : \mathcal{P}_s \rightarrow \mathbb{Z}[\alpha, t]$ be a linear mapping such that

1. $\psi(\alpha, 0)(P^n) = \alpha^n$;
2. One of the following equivalent conditions holds:

- (a) $\psi(\alpha, t_1)(e^{td}p) = \psi(\alpha, t + t_1)(p)$;
- (b) $\psi(\alpha, t)(dp) = \frac{\partial}{\partial t} \psi(\alpha, t)(p)$.

Then $\psi = f_1$.

Proof. Let conditions 1 and 2a hold. Then

$$\psi(\alpha, t + t_1)(p) = \psi(\alpha, t_1)(e^{td}p) = \psi(\alpha, 0)(e^{(t+t_1)d}p) = \xi_\alpha \Phi(t + t_1)p = f_1(\alpha, t_1 + t)(p).$$

It remains to prove that conditions 2a and 2b are equivalent.

Indeed, if $\psi(\alpha, t_1)(e^{td}p) = \psi(\alpha, t + t_1)(p)$, then $\psi(\alpha, t)(p) = \psi(\alpha, 0)(e^{td}p)$, therefore

$$\frac{\partial}{\partial t} \psi(\alpha, t)(p) = \frac{\partial}{\partial t} \psi(\alpha, 0)(e^{td}p) = \psi(\alpha, 0)(e^{td}dp) = \psi(\alpha, t)(dp).$$

Now let $\psi(\alpha, t)(dp) = \frac{\partial}{\partial t} \psi(\alpha, t)(p)$. Then

$$\frac{\partial}{\partial t} \psi(\alpha, t_1)(e^{td}p) = \psi(\alpha, t_1)(d(e^{td}p)) = \frac{\partial}{\partial t_1} \psi(\alpha, t_1)(e^{td}p)$$

Hence $\left(\frac{\partial}{\partial t} - \frac{\partial}{\partial t_1}\right) \psi(\alpha, t_1)(e^{td}p) = 0$, so $\psi(\alpha, t_1)(e^{td}p)$ depends on α and $t + t_1$.

Thus $\psi(\alpha, t_1)(e^{td}p) = \psi(\alpha, t + t_1)(e^{0d}p) = \psi(\alpha, t + t_1)(p)$. $\square$

Proposition 19 in the form 1, 2b was first proved in [Buch].

11 Ring $\mathcal{D}^*$

Let us identify $\mathcal{M} = \text{Qsym}[t_1, t_2, \dots]$. Let us remind that the representation $R : \mathcal{Z} \rightarrow \mathcal{D}$ is defined on the generators as $Z_k \rightarrow d_k$. Then R^* is the mapping

$$R^* : \mathcal{D}^*[\alpha] \rightarrow \text{Qsym}[t_1, t_2, \dots][\alpha] : (R^*\psi(\alpha)) Z^\omega = \psi(\alpha) (RZ^\omega).$$

The ring $\mathcal{D}^*[\alpha]$ is a graded ring with $\deg \alpha^i \psi_i = i + \deg \psi_i$.

Proposition 23.

$$R^* \varphi_{\alpha, P} = f(P).$$

For each m let us define the ring homomorphism $R_m^* : \mathcal{D}^*[\alpha] \rightarrow \text{Qsym}[t_1, \dots, t_m][\alpha]$:

$$R_m^*(\alpha, t_1, \dots, t_m)(\psi(\alpha)) = \psi(\alpha) \Phi(t_m) \dots \Phi(t_1) = R^*(\alpha, t_1, \dots, t_m, 0, 0, \dots).$$

Then $f_m(p) = R_m^* \varphi_{\alpha, p}$ for all $p \in \mathcal{P}$.

The restriction $R^* \rightarrow R_m$ is injective for the graduations $n \leq m$ and the mapping R^* is injective on the ring $\mathcal{D}^*[\alpha]$, since the mapping R is surjective.

Proposition 24. *Let $m \geq n$. Then the image of the n -th graded component of the ring $\mathcal{D}^*$ in the ring $\text{Qsym}[t_1, \dots, t_m]$ under the map R_m^* consists of all the homogeneous polynomials of degree n satisfying the relations:*

$$\begin{aligned} g(t_1, -t_1, t_3, \dots, t_m) &= g(0, 0, t_3, \dots, t_m); \\ g(t_1, t_2, -t_2, t_4, \dots, t_m) &= g(t_1, 0, 0, t_4, \dots, t_m); \\ &\dots \\ g(t_1, \dots, t_{m-2}, t_{m-1}, -t_{m-1}) &= g(t_1, \dots, t_{m-2}, 0, 0). \end{aligned}$$

Proof. The ring $\mathcal{D}^* \subset \mathcal{Z}^*$ consists of all the linear functions $\psi \in \mathcal{Z}^*$ satisfying the property:

$$\psi z_1 \Phi(t) \Phi(-t) z_2 = \psi z_1 z_2$$

for all $z_1, z_2 \in \mathcal{Z}$, that is the coefficients of all t^k , $k \geq 1$, on the left are equal to 0.

Since $R_m^* \psi(\alpha) = \psi(\alpha) \Phi(t_m) \dots \Phi(t_1)$, the relations of the theorem are valid.

On the other hand, let $g \in \text{Qsym}[t_1, \dots, t_m]$ be a homogeneous polynomial of degree n satisfying all the relations of the theorem. Let us prove that the corresponding linear function $\psi \in \mathcal{M}$ belongs to the image of $\mathcal{D}^*$ under the embedding R^* .

It is sufficient to prove the relation (12) in the case when z_1, z_2 are monomials. Since

$$\Phi(t) \Phi(-t) = 1 + (Z_1 - Z_1)t + (2Z_2 - Z_1^2)t^2 + \dots = 1 + (2Z_2 - Z_1^2)t^2 + \dots, \text{ and } \deg \psi = n,$$

the cases $\deg z_1 + \deg z_2 = n$ and $n - 1$ are trivial. Let

$$z_1 = Z^\omega, \quad z_2 = Z^{\omega'}, \quad \omega = (j_1, \dots, j_l), \quad \omega' = (j'_1, \dots, j'_{l'}), \quad |\omega| + |\omega'| = n - k, \quad k \geq 2.$$

Then the only equality we need to prove is

$$\psi Z^\omega \left(\sum_{i=0}^k (-1)^i Z_{k-i} Z_i \right) Z^{\omega'} = 0$$

Let us consider the equality

$$g(t_1, \dots, t_l, t_{l+1}, -t_{l+1}, t_{l+3}, \dots, t_{l+2+l'}, \dots, t_m) = g(t_1, \dots, t_l, 0, 0, t_{l+3}, \dots, t_{l+2+l'}, \dots, t_m).$$

The coefficient of the monomial $t_1^{j_1} \dots t_l^{j_l} t_{l+1}^{j_{l+1}} t_{l+3}^{j'_{l+3}} \dots t_{l+2+l'}^{j'_{l+2+l'}}$ on the left is exactly

$$\sum_{i=0}^k (-1)^i \psi Z^\omega Z_{k-i} Z_i Z^{\omega'} = \psi Z^\omega \left(\sum_{i=0}^k (-1)^i Z_{k-i} Z_i \right) Z^{\omega'},$$

and on the right it is equal to 0. So $\psi Z^\omega \Phi(t) \Phi(-t) Z^{\omega'} = 0$ for all ω, ω' , therefore $\psi \in \mathcal{D}^*$. $\square$

Proposition 25. *The image of the space $\mathcal{P}^n$ under the mapping $\varphi : \mathcal{P} \rightarrow \mathcal{D}^*[\alpha]$ consists of all homogeneous functions $\psi(\alpha)$ of degree n such that*

$$\psi(-\alpha)\Phi(\alpha)D = \psi(\alpha)D \quad (12)$$

for all $D \in \mathcal{D}$.

Proof. According to Proposition 3 we have $\xi_{-\alpha}\Phi(\alpha) = \xi_\alpha$, so if $\psi(\alpha) = \varphi_{\alpha,p}$, then

$$\psi(-\alpha)\Phi(\alpha)D = \xi_{-\alpha}\Phi(\alpha)Dp = \xi_\alpha Dp = \psi(\alpha)D.$$

On the other hand, let condition (12) hold. Then the polynomial $g(\alpha, t_1, \dots, t_n) = R_n^*\psi(\alpha)$ satisfies the relation

$$g(-\alpha, t_1, \dots, t_{n-1}, \alpha) = \psi(-\alpha)\Phi(\alpha)\Phi(t_{n-1}) \dots \Phi(t_1) = \psi(\alpha)\Phi(0)\Phi(t_{n-1}) \dots \Phi(t_1) = g(\alpha, t_1, \dots, t_{n-1}, 0).$$

Proposition 24 implies that the polynomial g satisfies all the relations of Theorem 2, so $g = f_n(p^n)$. We have $R_n^*\psi(\alpha) = g = f_n(p^n) = R_n^*\varphi_{\alpha,p^n}$. Since the restriction $R^* \rightarrow R_n^*$ is injective on j -th graded component of the ring $\mathcal{D}^*$, $j \leq n$, and R^* is an embedding, we obtain that $\psi(\alpha) = \varphi_{\alpha,p^n}$. $\square$

There is a right action of the ring $\mathcal{D}$ on it's graded dual $\mathcal{D}^*$:

$$(\psi, D) \rightarrow \psi D : \quad (\psi D)(D') = \psi(DD').$$

If $\deg \psi = n$ and $\deg D = k$, then $\deg \psi D = n - k$. Similarly, there is an action of $\mathcal{D}[[\alpha]]$ on $\mathcal{D}^*[[\alpha]]$. Then condition (12) means that $\psi(-\alpha)\Phi(\alpha) = \psi(\alpha)$. Since for any homogeneous function $\psi(\alpha)$ of degree n the function $\psi(\alpha)\Phi(\alpha)$ still has degree n , we obtain the corollary.

Corollary 26. *Let $\psi(\alpha) \in \mathcal{D}^*[\alpha]$. Then*

$$\psi(\alpha) \in \varphi(\mathcal{P}) \Leftrightarrow \psi(-\alpha)\Phi(\alpha) = \psi(\alpha). \quad (13)$$

Let $\psi(\alpha) = \psi_0 + \psi_1\alpha + \dots + \psi_n\alpha^n$, $u(\alpha) = \psi_0 + \psi_2\alpha^2 + \psi_4\alpha^4 + \dots$, $v = \psi_1\alpha + \psi_3\alpha^3 + \dots$. Then $u(\alpha)\Phi(\alpha) - v(\alpha)\Phi(\alpha) = u(\alpha) + v(\alpha)$. Therefore

$$u(\alpha)(\Phi(\alpha) - 1) = v(\alpha)(\Phi(\alpha) + 1). \quad (14)$$

For example,

$$\psi_0 d = 2\psi_1; \quad \psi_0 d_3 + \psi_2 d = \psi_1 d_2 + 2\psi_3.$$

Consider the ring $\mathcal{D}^* \otimes \mathbb{Z}[\frac{1}{2}]$. Then for the graded dual Hopf algebra we have: $(\mathcal{D} \otimes \mathbb{Z}[\frac{1}{2}])^* \simeq \mathcal{D}^* \otimes \mathbb{Z}[\frac{1}{2}]$. The condition that describes the image of the ring $\mathcal{P} \otimes \mathbb{Z}[\frac{1}{2}]$ in $\mathcal{D}^* \otimes \mathbb{Z}[\frac{1}{2}][\alpha]$ is the same, namely relation (14).

Proposition 27. *In the ring $\mathcal{D}^* \otimes \mathbb{Z}[\frac{1}{2}][\alpha]$ relation (14) is equivalent to the relation:*

$$v(\alpha) = u(\alpha) \frac{\Phi(\alpha) - 1}{\Phi(\alpha) + 1} = u(\alpha) \frac{\frac{\Phi(\alpha) - 1}{2}}{1 + \frac{\Phi(\alpha) - 1}{2}} = u(\alpha) \sum_{k=1}^{\infty} (-1)^{k-1} \left(\frac{\Phi(\alpha) - 1}{2} \right)^k. \quad (15)$$

Proof. Of course, relation (15) follows from relation (14). On the other hand, we need to prove, that the function $v(\alpha)$ defined by relation (14) is odd, if $u(\alpha)$ is even. Indeed,

$$\begin{aligned} v(-\alpha) &= u(-\alpha) \frac{\Phi(-\alpha) - 1}{\Phi(-\alpha) + 1} = u(\alpha) \frac{\frac{1}{\Phi(\alpha)} - 1}{\frac{1}{\Phi(\alpha)} + 1} = u(\alpha)(1 - \Phi(\alpha))\Phi(\alpha)^{-1}(1 + \Phi(\alpha))^{-1}\Phi(\alpha) = \\ &= -u(\alpha) \frac{\Phi(\alpha) - 1}{\Phi(\alpha) + 1} = -v(\alpha). \end{aligned}$$

$\square$

Let us take $\varphi_{\alpha,p}$ with $\alpha = 0$. Then we obtain the classical map

$$\varphi_0 : \mathcal{P} \rightarrow \mathcal{D}^* : p \rightarrow \varphi_{0,p} \quad \varphi_0(p)D = \xi_0 Dp, \quad \forall p \in \mathcal{P}, D \in \mathcal{D}$$

Proposition 28. *The mapping $\varphi_0 \otimes 1 : \mathcal{P} \otimes \mathbb{Z}[\frac{1}{2}] \rightarrow \mathcal{D}^* \otimes \mathbb{Z}[\frac{1}{2}]$ is a surjection.*

Proof. Let $\psi_0 \in \mathcal{D}^* \otimes \mathbb{Z}[\frac{1}{2}]$. Consider the element $\psi(\alpha) = u(\alpha) + v(\alpha)$, $u(\alpha) = \psi_0$, $v(\alpha) = \psi_0 \frac{\Phi(\alpha)-1}{\Phi(\alpha)+1}$. According to $\mathbb{Z}[\frac{1}{2}]$ -version of Corollary 26 we obtain that $\psi(\alpha) = \varphi_{\alpha,p} \in \varphi \otimes 1(\mathcal{P} \otimes \mathbb{Z}[\frac{1}{2}])$. Then $\psi_0 = \psi(0) = \varphi_{0,p}$. $\square$

Question 1. What is the image of the map φ_0 over the integers?

Remark. Let us note, that the space $\mathcal{P}/\text{Ker } \varphi$ consists of the equivalence classes of integer combinations of polytopes under the equivalence relation: $p \sim q$ if and only if p and q have equal flag f-vectors. Rank of the n -th graded component of the group $\mathcal{P}/\text{Ker } \varphi$ is equal to c_n ([BB], see also Section 6).

On the other hand, by Corollary 14 rank of the n -th graded component of the ring $\mathcal{D}^*$ is equal to c_{n-1} , so the mapping $\varphi_0 : \mathcal{P}/\text{Ker } \varphi \rightarrow \mathcal{D}^*$ is not injective.

Example. Let us consider small dimensions.

- $n = 1$. $\mathcal{P}/\text{Ker } \varphi$ is generated by $CC = I$. The ring $\mathcal{D}$ in this graduation is generated by d . Then $\varphi_0(I) = 2d^*$.
- $n = 2$. $\mathcal{P}/\text{Ker } \varphi$ is generated by $CCC = \Delta^2$ and $BCC = I^2$. $\mathcal{D}$ has one generator d_2 . Then $\varphi_0(\Delta^2) = 3d_2^*$, $\varphi_0(I^2) = 4d_2^*$. Therefore $d_2^* = \varphi_0(I^2 - \Delta^2)$, $\text{Ker } \varphi_0$ is generated by $3I^2 - 4\Delta^2$.
- $n = 3$. $\mathcal{P}/\text{Ker } \varphi$ is generated by $BCCC, CBCC = CI^2, CCCC = \Delta^3$, while $\mathcal{D}$ is generated by d_3, d_2d . Then

$$\varphi_0(BC^3) = 5d_3^* + 18(d_2d)^*, \quad \varphi_0(CBC^2) = 5d_3^* + 16(d_2d)^*, \quad \varphi_0(C^4) = 4d_3^* + 12(d_2d)^*$$

So $\text{Im } \varphi_0$ has the basis $d_3^*, 2(d_2d)^*$ and $\text{Ker } \varphi_0$ is generated by $2BC^3 - 6CBC^2 + 5C^4$.

Here by D_ω^* we denote the element of the dual basis: $\langle D_\omega^*, D_\sigma \rangle = \delta_{\omega,\sigma}$

12 Multiplicative Structure

Theorem 4. *The ring $f(\mathcal{P}) \otimes \mathbb{Q}$ is a free polynomial algebra.*

Proof. According to Proposition 17 and Corollary 7 to the shuffle algebra structure theorem we have:

$$\mathcal{D}^* \otimes \mathbb{Q} \simeq \mathbb{Q}[\text{LYN}_{\text{odd}}],$$

Where LYN_{odd} are Lyndon words consisting of odd positive integers. This is a free polynomial algebra. Therefore $\mathcal{D}^* \otimes \mathbb{Q}[\alpha]$ is a free polynomial algebra in the generators LYN_{odd} and α .

Let $\{f_\lambda\}$ be the elements of $\mathcal{D}^* \otimes \mathbb{Q}$ corresponding to the Lyndon words. Consider the functions $f_\lambda(\alpha)$ defined as

$$f_\lambda(\alpha) = u_\lambda(\alpha) + v_\lambda(\alpha); \quad u_\lambda(\alpha) = \psi_\lambda, \quad v_\lambda(\alpha) = u_\lambda(\alpha) \frac{\Phi(\alpha) - 1}{\Phi(\alpha) + 1} = \psi_\lambda \frac{e^{s(\alpha)} - 1}{e^{s(\alpha)} + 1}$$

Let us identify $\mathcal{D}^*$ with its image in $\text{Qsym}[t_1, t_2, \dots]$ under the embedding R^* .

Then $f_\lambda(\alpha) \in f(\mathcal{P} \otimes \mathbb{Q}) = f(\mathcal{P}) \otimes \mathbb{Q}$.

Lemma 4. $f(\mathcal{P}) \otimes \mathbb{Q} = \mathbb{Q}[f_\lambda, \alpha^2]$.

Proof. At first let us proof that the polynomials $\{f_\lambda\}, \alpha^2$ are algebraically independent.

Let $g \in \mathbb{Q}[x, y_1, y_2, \dots, y_s]$ be a polynomial such that $g(\alpha^2, f_1(\alpha), \dots, f_s(\alpha)) = 0$. We can write this equality as

$$g_0(f_1(\alpha), \dots, f_s(\alpha)) + g_2(f_1(\alpha), \dots, f_s(\alpha))\alpha^2 + \dots + g_{2t}(f_1(\alpha), \dots, f_s(\alpha))\alpha^{2t} = 0$$

for some integers s, t . Let us set $\alpha = 0$. Then we obtain

$$g_0(f_1, \dots, f_s) = 0$$

Since $\{f_\lambda\}$ are algebraically independent, $g_0 \equiv 0$. Let us divide the equality by α^2 to obtain

$$g_2(f_1(\alpha), \dots, f_s(\alpha)) + g_2(f_1(\alpha), \dots, f_s(\alpha))\alpha^2 + \dots + g_{2t}(f_1(\alpha), \dots, f_s(\alpha))\alpha^{2t-2} = 0$$

Iterating this step in the end we obtain that all the polynomials $g_{2i} \equiv 0$, $i = 0, 1, \dots, 2t$, so $g \equiv 0$.

Now let us prove that the polynomials $\{f_\lambda(\alpha)\}, \alpha^2$ generate $f(\mathcal{P}) \otimes \mathbb{Q}$.

Let $\psi(\alpha) = \psi_0 + \psi_1\alpha + \dots + \psi_n\alpha^n \in f(\mathcal{P}) \otimes \mathbb{Q}$. The polynomials $\{f_\lambda\}$ generate $\mathcal{D}^*$, so we can find a polynomial $g_0 \in \mathbb{Q}[y_1, \dots, y_s]$ such that

$$g_0(f_1, \dots, f_s) = \psi_0$$

Let us consider the element $\psi(\alpha) - g_0(f_1(\alpha), \dots, f_s(\alpha)) \in f(\mathcal{P}) \otimes \mathbb{Q}$. Since $\theta_1 = \theta_0 \frac{d}{d\alpha}$ for any

$$\theta(\alpha) = \theta_0 + \theta_1\alpha + \theta_2\alpha^2 + \dots + \theta_r\alpha^r \in f(\mathcal{P}) \otimes \mathbb{Q},$$

we obtain that

$$\psi(\alpha) - g_0(f_1(\alpha), \dots, f_s(\alpha)) = \hat{\psi}(\alpha)\alpha^2.$$

$\hat{\psi}(\alpha)\alpha^2 \in f(\mathcal{P}) \otimes \mathbb{Q}$, so $\hat{\psi}(-\alpha)\alpha^2\Phi(\alpha) = \hat{\psi}(\alpha)\alpha^2$. Since $\mathcal{D}^* \otimes \mathbb{Q}[\alpha]$ is a polynomial ring, $\hat{\psi}(-\alpha)\Phi(\alpha) = \hat{\psi}(\alpha)$, so $\hat{\psi}(\alpha) \in f(\mathcal{P}) \otimes \mathbb{Q}$.

Iterating this argument in the end we obtain the expression of $\psi(\alpha)$ as a polynomial in $\{f_\lambda\}$ and α^2 . $\square$

This lemma proves the theorem. $\square$

Let us note that dimension of the n -th graded component of the ring $f(\mathcal{P}) \otimes \mathbb{Q}$ is equal to the n -th Fibonacci number c_n .

Corollary 29. *There is an isomorphism of free polynomial algebras*

$$f(\mathcal{P}) \otimes \mathbb{Q} \simeq \mathbb{Q}[\text{LYN}_{odd}^n, \alpha^2] = \mathcal{N}_{odd} \otimes \mathbb{Q}[\alpha^2] \simeq \mathbb{Q}[\text{LYN}_{12}^n] = \mathcal{N}_{12} \otimes \mathbb{Q}.$$

Proof. Indeed, these three algebras are free polynomial algebras with the same dimensions of the corresponding graded components: dimension of the n -th graded component of each algebra is equal to c_n . $\square$

Let us denote by k_n the number of multiplicative generators of degree n . According to Corollary 29 for $n \geq 3$ the number of generators k_n is equal to the number of Lyndon words in LYN_{odd} or in LYN_{12} of degree n .

For example, for small n we have

n	LYN_{12}^n	LYN_{odd}^n
3	[1,2]	[3]
4	[1,1,2]	[1,3]
5	[1,2,2], [1,1,1,2]	[5], [1,1,3]
6	[1,1,2,2], [1,1,1,1,2]	[1,5], [1,1,1,3]
7	[1,2,2,2], [1,1,2,1,2], [1,1,1,2,2], [1,1,1,1,1,2]	[7], [1,1,5], [1,3,3], [1,1,1,1,3],

Corollary 30. 1. $k_{n+1} \geq k_n$

2. $k_n \geq N_n - 2$, where N_n is the number of different decompositions of n into the sum of odd numbers.

Remark. It is a well-known fact, that the number of the decompositions of n into the sum of odd numbers is equal to the number of the decompositions of n into the sum of different numbers. For example,

$$\begin{array}{rclcl} 2 & = & 1 + 1 & = & 2 \\ 3 & = & 1 + 1 + 1, 3 & = & 1 + 2, 3 \\ 4 & = & 1 + 1 + 1 + 1, 1 + 3 & = & 1 + 3, 4 \\ 5 & = & 1 + 1 + 1 + 1 + 1, 1 + 1 + 3, 5 & = & 1 + 4, 2 + 3, 5 \end{array}$$

Proof. 1. $k_1 = k_2 = k_3 = 1$.

Let $n \geq 3$ and let $w = [a_1, \dots, a_k] \in \text{LYN}_{12}^n$. Then $1w = [1, a_1, \dots, a_k] \in \text{LYN}_{12}^{n+1}$. Indeed, for any proper tail $[a_i, \dots, a_k]$, $i \geq 1$ if $a_i > 1$, then $[a_i, \dots, a_k] > 1w$. If $a_i = 1$ then $i \neq k$ and $[a_{i+1}, \dots, a_k] > [a_1, \dots, a_k]$, since w is Lyndon. Thus $k_{n+1} \geq k_n$.

2. Any decomposition $n = d_1 + \dots + d_k$ into the sum of odd numbers $d_1 \leq d_2 \leq \dots \leq d_k$ gives a Lyndon word $[d_1, \dots, d_k] \in \text{LYN}_{\text{odd}}$, except for the case, when $d_1 = \dots = d_k = d \neq n$.

If $d \geq 5$, then the word $[1, d-2, 1, d, \dots, d]$ is Lyndon. When $d = 1$ or $d = 3$ there can be no Lyndon word corresponding to the decomposition $n = 1 + \dots + 1$ or $n = 3 + \dots + 3$, as it happens for $n = 6$:

$$6 = 1 + 5 = 3 + 3 = 1 + 1 + 1 + 3 = 1 + 1 + 1 + 1 + 1$$

We have $k_6 = 2$ and $N_6 = 4$. Thus $k_n \geq N_n - 2$. □

Then we obtain the corollary:

Corollary 31. For any t such that $|t| < \frac{\sqrt{5}-1}{2}$ there exists a decomposition of the generating series of Fibonacci numbers as an infinite product:

$$\frac{1}{1-t-t^2} = 1 + t + t^2 + 2t^3 + 3t^4 + \dots = \sum_{n=0}^{\infty} c_n t^n = \prod_{i=1}^{\infty} \frac{1}{(1-t^i)^{k_i}}$$

Where the infinite product $\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{1}{(1-t^i)^{k_i}}$ converges absolutely.

The numbers k_n satisfy the properties $k_{n+1} \geq k_n \geq N_n - 2$, where N_n is the number of the decompositions of n into the sum of odd summands.

Proof. Let $0 \leq t < \frac{\sqrt{5}-1}{2}$. Then the series $\sum_{n=0}^{\infty} c_n t^n$ converges. This means that $R_N = \sum_{n=N}^{\infty} c_n t^n \xrightarrow{N \rightarrow \infty} 0$.

But

$$S_n = \sum_{k=0}^{\infty} c_k t^k - \prod_{i=1}^n \frac{1}{(1-t^i)^{k_i}} = a_{n+1} t^{n+1} + a_{n+2} t^{n+2} + \dots$$

where $0 \leq a_i \leq c_i$. Then $0 \leq S_n \leq R_n$. Therefore $S_n \xrightarrow{n \rightarrow \infty} 0$.

On the other hand, there is a criterion of an absolute convergence of an infinite product:

The product $\prod_{n=1}^{\infty} (1 + b_n)$ converges absolutely if and only if the series $\sum_{n=1}^{\infty} b_n$ converges absolutely.

It follows from the fact that $\frac{|\ln(1+b_n)|}{|b_n|} = \left| \frac{\ln(1+b_n)}{b_n} \right| \xrightarrow{n \rightarrow \infty} 1$, when $b_n \xrightarrow{n \rightarrow \infty} 0$.

In our case the series $-\sum_{i=1}^{\infty} k_i \ln(1-t^i)$ converges for all $0 \leq t < \frac{\sqrt{5}-1}{2}$, therefore $\sum_{i=1}^{\infty} k_i t^i$ converges for the same values of t . Therefore radius of convergence of this series is at least $\frac{\sqrt{5}-1}{2}$. This implies that the product converges absolutely for $|t| < \frac{\sqrt{5}-1}{2}$. □

Here are the examples in small dimensions:

n	generators	additive basis	c_n	k_n
1	[1]	[1]	1	1
2	[2]	[2], [1]*[1]	2	1
3	[1,2]	[1,2], [2]*[1], [1]*[1]*[1]	3	1
4	[1,1,2]	[1,1,2], [1,2]*[1], [2]*[2], [2]*[1]*[1], [1]*[1]*[1]*[1]*[1]	5	1
5	[1,2,2], [1,1,1,2]	[1,2,2], [1,1,1,2], [1,1,2]*[1], [1,2]*[2], [1,2]*[1]*[1]	8	2
6	[1,1,2,2], [1,1,1,1,2]	[2]*[2]*[1], [2]*[1]*[1]*[1], [1]*[1]*[1]*[1]*[1]	13	2
7	[1,2,2,2], [1,1,2,1,2], [1,1,1,2,2], [1,1,1,1,1,2]		21	4

The corresponding products are:

$$\begin{aligned}
(1-t)(1-t^2) &= 1-t-t^2+t^3; \\
(1-t)(1-t^2)(1-t^3) &= 1-t-t^2+t^4+t^5-t^6; \\
(1-t)(1-t^2)(1-t^3)(1-t^4) &= 1-t-t^2+2t^5-t^8-t^9+t^{10}; \\
(1-t)(1-t^2)(1-t^3)(1-t^4)(1-t^5)^2 &= 1-t-t^2+2t^6+2t^7-t^8-t^9-2t^{10}- \\
&\quad -t^{11}-t^{12}+2t^{13}+2t^{14}-t^{18}-t^{19}+t^{20}.
\end{aligned}$$

The numbers k_i can be found in the following way:

$$\begin{aligned}
-\log(1-t-t^2) &= -\sum_{i=1}^{\infty} k_i \log(1-t^i); \\
\sum_{n=1}^{\infty} \frac{1}{n} \sum_{j=0}^n \binom{n}{j} t^{n+j} &= \sum_{i=1}^{\infty} k_i \sum_{r=1}^{\infty} \frac{t^{ri}}{r}
\end{aligned}$$

Thus for any N

$$\sum_{j=0}^{\lfloor \frac{N}{2} \rfloor} \frac{\binom{N-j}{j}}{N-j} = \frac{1}{N} \sum_{i|N} i k_i.$$

According to the Möbius inversion formula we obtain

$$N k_N = \sum_{d|N} d \sum_{j=0}^{\lfloor \frac{d}{2} \rfloor} \frac{\binom{d-j}{j}}{d-j} \mu\left(\frac{N}{d}\right),$$

Where $\mu(n)$ is the Möbius function, that is

$$\mu(n) = \begin{cases} 1, & n = 1; \\ (-1)^r, & n = p_1 \dots p_r, \{p_i\} - \text{distinct prime numbers}; \\ 0, & n \text{ is not square-free.} \end{cases}$$

For example, if N is a prime number $N = p$, then

$$p k_p = -1 + p \sum_{j=0}^{\lfloor \frac{p}{2} \rfloor} \frac{\binom{p-j}{j}}{p-j}; \quad k_p = \sum_{j=1}^{\lfloor \frac{p}{2} \rfloor} \frac{\binom{p-j}{j}}{p-j}.$$

Each summand is integer, since $\binom{p-j}{j} = \frac{\binom{p-j-1}{j-1}(p-j)}{j}$ is integer, and $(p-j)$ and j are relatively prime numbers.

Thus $k_5 = 2$, $k_7 = 4$, $k_{11} = 18$, and so on.

13 Bayer-Billera Ring

Let us consider the free graded abelian group $BB \subset \mathcal{P}$, generated by 1 and all the polytopes $Q \in \Omega^n$, $n = 1, 2, \dots$. Rank of the n -th graded component of this group is equal to c_n .

Since the determinant of the matrix K^n is equal to 1, the generalized f -polynomials $\{f(Q), Q \in \Omega^n\}$ form a basis of the n -th graded component of the ring $f(\mathcal{P})$.

So the composition of the inclusion $i : BB \subset \mathcal{P}$ and the mapping $f : \mathcal{P} \rightarrow \text{Qsym}[t_1, t_2, \dots][\alpha]$ is an isomorphism of the abelian groups BB and $f(\mathcal{P})$. This gives a projection $\pi : \mathcal{P} \rightarrow BB$

$$\pi(p) = x \in BB : f(p) = f(x).$$

It follows from the definition, that $\pi \circ i = 1$ on the space BB .

Theorem 5. *The projection $\pi : \mathcal{P} \rightarrow BB$ defined by the relation $f(\pi p) = f(p)$ gives the group BB the structure of a commutative associative ring with the multiplication $x * y = \pi(x \times y)$, such that $f(x * y) = f(x)f(y)$ and $BB \otimes \mathbb{Q}$ is a free polynomial algebra in a countable set of variables.*

14 Rota-Hopf Algebra

14.1 Hopf Algebra of Graded Posets

Definition 23. Let P be a finite poset with a minimal element $\hat{0}$ and a maximal element $\hat{1}$. An element y in P covers another element x in P , if $x < y$ and there is no z in P such that $x < z < y$. A poset P is called *graded*, if there exists a rank function $\rho : \mathcal{P} \rightarrow \mathbb{Z}$ such that $\rho(\hat{0}) = 0$ and $\rho(x) + 1 = \rho(y)$ if y covers x . Let us denote $\rho(x, y) = \rho(y) - \rho(x)$ for $x \leq y$, and $\rho(P) = \rho(\hat{1})$. Two finite graded posets are isomorphic if there exists an order preserving bijection between them.

Let $\mathcal{R}$ be the vector space over the field k with basis the set of all isomorphism classes of finite graded posets. $\mathcal{R}$ is a graded connected Hopf algebra where the degree of P is its rank $\rho(P)$, the multiplication is a cartesian product of posets

$$P \cdot Q = P \times Q, \quad (x, y)_{P \times Q} \leq (u, v) \text{ if and only if } x \leq u \text{ and } y \leq v,$$

the unit element is the poset with one element $\hat{0} = \hat{1}$, the comultiplication is

$$\Delta(P) = \sum_{\hat{0} \leq z \leq \hat{1}} [\hat{0}, z] \otimes [z, \hat{1}],$$

where $[x, y]$ is the subposet $\{z \in P | x \leq z \leq y\}$, and the counit

$$\varepsilon(P) = \begin{cases} 1, & \text{if } \hat{0} = \hat{1}; \\ 0, & \text{else} \end{cases}$$

The antipode of the algebra $\mathcal{R}$ is given by the formula [Sch1]:

$$S(P) = \sum_{k \geq 0} \sum_{\hat{0} = x_0 < x_1 < \dots < x_k = \hat{1}} (-1)^k [x_0, x_1] \cdot [x_1, x_2] \dots [x_{k-1}, x_k]$$

Example. For the simplest Boolean algebra $B_1 = \{\hat{0}, \hat{1}\}$, which is the face lattice of the point pt

$$\begin{aligned} \Delta(B_1) &= 1 \otimes B_1 + B_1 \otimes 1 \\ S(B_1) &= -B_1 \end{aligned}$$

The Hopf algebra of graded posets was originated in the work by Joni and Rota [JR]. Variations of this construction were studied in [Ehr, ABS, Sch1, Sch2]. A generalization of this algebra can be found in [RS].

14.2 Natural Homomorphism from the Ring of Polytopes to the Rota-Hopf algebra

There is a natural linear mapping $L : \mathcal{P} \rightarrow \mathcal{R}$ of degree +1 that sends a polytope P to its face lattice $L(P)$.

It is easy to see that this mapping is injective, but it is not a ring homomorphism, since it doesn't preserve a unit: it sends the point pt to the Boolean algebra $L(\text{pt}) = B_1$, while the unit of $\mathcal{R}$ is a one-element set $\{\hat{0} = \hat{1}\}$.

Remark. Let us note that $L(P \times Q) \neq L(P) \times L(Q)$, since the face lattice $L(P \times Q)$ contains the empty face $\emptyset$, which can be considered as $\emptyset \times \emptyset$, but evidently has no faces of the form $F \times \emptyset$ or $\emptyset \times G$, where F and G are non-empty faces of P and Q respectively.

Definition 24. There is a natural linear mapping $l = L * d* : \mathcal{P} \rightarrow \mathcal{R}$ of degree 0 that sends the polytope to the sum of the face lattices of its vertex figures (Let us recall that the linear operator $*$ sends the polytope P to its polar P^* .)

$$l(\text{pt}) = \{\hat{0}\}, \quad l(P) = L((dP^*)^*) = \sum_{v-\text{vertex}} L(P/v).$$

Example. To illustrate the mapping l let us consider the sets of simplicial and simple polytopes:

- For a simplicial polytope P^n each vertex figure is again a simplicial polytope. Combinatorially we have $\partial(P^n/v) = \text{link}_{\partial P^n}(v)$. Therefore the operation $*d*$ preserves the linear space of all simplicial polytopes.
- For a simple polytope P^n all of its vertex figures are simplices, therefore $*d*(P^n) = f_0(P^n)\Delta^{n-1}$ and $l(P^n) = f_0 L(\Delta^{n-1}) = f_0 \{\hat{0}, \hat{1}\}^n = f_0 B_1^n = B_n$, where $f_0(P^n)$ is the number of vertices of P^n , and $\{\hat{0}, \hat{1}\}^n = B_1^n = B_n$ is a Boolean algebra. For example, $l(\Delta^n) = (n+1)B_1^n$ and $L(\Delta^n) = B_1^{n+1}$.

Proposition 32. l is a homomorphism of graded rings.

Proof. We have

$$l(P) = \sum_{v-\text{vertex}} L(P/v) = \sum_v [v, P],$$

where L/v is a vertex figure, and $[v, P]$ is the interval between the vertex v and the polytope P in the face lattice $L(P)$. Then

$$l(P \times Q) = \sum_{v \times w} [v \times w, P \times Q] = \sum_{v \times w} [v, P] \times [w, Q] = \left(\sum_v [v, P] \right) \cdot \left(\sum_w [w, Q] \right) = l(P) \cdot l(Q)$$

Here v, w are vertexes of P and Q respectively. □

Definition 25. Consider the linear span of all Boolean algebras $B_n = \{0, 1\}^n = L(\Delta^{n-1})$, $n \geq 0$

$$\mathcal{B} = \text{Ls}(1, B_1, B_2, \dots) \subset \mathcal{R}.$$

We have $B_i B_j = B_{i+j}$, so it is a subring in $\mathcal{R}$. Let us denote $x = B_1$. Since $\Delta x = 1 \otimes x + x \otimes 1$, it is a Hopf subalgebra isomorphic to the Hopf algebra $\mathbb{k}[x]$, $\Delta x = 1 \otimes x + x \otimes 1$.

Proposition 33. The image of the ring of simple polytopes $\mathcal{P}_s$ under the mapping l is a $\mathbb{Z}$ -subalgebra in $\mathcal{B}$ multiplicatively generated by $2x$ and x^2 , that is

$$l(\mathcal{P}_s) = \mathbb{Z}[x_1, x_2] / (x_1^2 - 4x_2), \quad \deg x_1 = 1, \deg x_2 = 2$$

where x_1 corresponds to $2x = 2B_1$ and x_2 to $x^2 = B_2$

Proof. Indeed, for a simple polytope P^n we have $l(P^n) = f_0 x^n$. On the other hand, we know that for any simple polytope $2f_1 = n f_0$, therefore for odd n the number of vertices f_0 should be even. So $l(\mathcal{P}_s) \in \mathbb{Z}[2x, x^2]$. On the other hand, $2x = l(I)$ and $x^2 = l(I^2 - \Delta^2)$. □

In [Ehr] Richard Ehrenborg introduced the F -quasi-symmetric function of a graded poset of rank n

$$F(P) = \sum_{\hat{0}=x_0 < x_1 < \dots < x_{k+1}=\hat{1}} M_{(\rho(x_0, x_1), \rho(x_1, x_2), \dots, \rho(x_k, x_{k+1}))} = \sum_{0 < a_1 < \dots < a_k < n} f_{a_1, \dots, a_k}(P) M_{(a_1, a_2 - a_1, \dots, n - a_k)}$$

where the sum ranges over all chains from $\hat{0}$ to $\hat{1}$, and $f_{a_1, \dots, a_k}$ are flag numbers. This mapping induces a Hopf algebra homomorphism $F : \mathcal{R} \rightarrow \text{Qsym}[t_1, t_2, \dots]$.

Proposition 34. *For a polytope P^n the following relation holds:*

$$F(l(P^n))^* = f(0, t_1, t_2, \dots)(P^n)$$

Thus

$$* \circ F \circ l = f|_{\alpha=0}.$$

Proof. For the point pt the relation is trivial $F(l(\text{pt}))^* = 1 = f(0, t_1, t_2, \dots)(\text{pt})$.

Since in the poset $L(P/v) = [v, P]$ rank of v is equal to its dimension 0, $\rho(F) = \dim F$, and we have:

$$\begin{aligned} F(l(P^n))^* &= \sum_v F([v, P^n])^* = \sum_{v \subset F^{a_2} \subset \dots \subset F^{a_k} \subset P^n} M_{(n-a_k, \dots, a_2)} = \\ &= \sum_{0 < a_2 < \dots < a_k < n} f_{0, a_2, \dots, a_k} \alpha^0 M_{(n-a_k, \dots, a_2-0)} = f(0, t_1, t_2, \dots)(P^n) \end{aligned}$$

□

Proposition 35. *We have*

$$L = l \circ (C - \frac{1}{2}[I]),$$

where C is a cone operator and $[I]$ is the multiplication by the interval I . So the images of the maps L and l over $\mathbb{Z}[\frac{1}{2}]$ in dimensions $n \geq 1$ coincide.

Proof. Let us mention that $1 = \{\hat{0}\}$ does not belong to the image of L .

For a polytope P we have $L(CP) = \{\hat{0}, \hat{1}\} \cdot L(P) = L(\text{pt}) \cdot L(P)$.

On the other hand, $l(I) = 2\{\hat{0}, \hat{1}\} = 2L(\text{pt})$. So $L(\text{pt})$ belongs to the image of l if and only if 2 is invertable.

Let P be an n -dimensional polytope, $n \geq 1$. Then

$$\begin{aligned} l(CP) &= L(*d * CP) = L(*dCP^*) = L(*(P^* + CdP^*)) = L(P) + L(C * d * P) = \\ &= L(P) + L(\text{pt}) \cdot l(P) = L(P) + \frac{1}{2}l(I) \cdot l(P) = L(P) + l\left(\frac{1}{2}I \times P\right) \end{aligned}$$

So $L(P) = l(CP - \frac{1}{2}I \times P)$. □

Remark. The homomorphism l is not injective: we see that on the ring of simple polytopes it remembers only the number of vertices. However Proposition 35 shows that l is injective on the image of the operator $C - \frac{1}{2}[I]$.

Corollary 36.

$$F(L(P))^* = Cf(P)|_{\alpha=0} - \sigma_1 f(P)|_{\alpha=0},$$

where $\sigma_1 = \sum_{i=1}^{\infty} t_i = M_{(1)}$.

Proof. This formula follows directly from Proposition 35. □

Let us denote the linear mapping $\text{Qsym}[t_1, t_2, \dots][\alpha] \xrightarrow{(C-\sigma_1)|_{\alpha=0}} \text{Qsym}[t_1, t_2, \dots]$ by Λ . Then on elementary monomials Λ has a very simple form

$$\Lambda(\alpha^{a_1} M_{(n-a_k, \dots, a_2-a_1)}) = M_{(n-a_k, \dots, a_2-a_1, a_1+1)}$$

Example. For a simple polytope P^n we have $l(P^n) = f_0 \cdot L(\Delta^{n-1}) = f_0 x^n$,

$$F(l(P^n)) = F(l(P^n))^* = f_0 \sigma_1^n = f(0, t_1, t_2, \dots)(P^n) = f_1(0, \sigma_1)(P^n).$$

Using Corollary 36 we obtain that

$$\begin{aligned} F(L(P^n))^* &= (C - \sigma_1) f_1(\alpha, \sigma_1)(P^n)|_{\alpha=0} = \sum_{i=1}^{\infty} t_i f_1(t_i, t_1 + \dots + t_{i-1})(P^n) = \\ &= \sum_{j=1}^{\infty} t_j^{n+1} + \sum_{i=1}^n f_{n-i} \left(\sum_{j=1}^{\infty} t_j^{n+1-i} (t_1 + \dots + t_{j-1})^i \right) = \\ &= M_{(n+1)} + \sum_{i=1}^n f_{n-i} \sum_{i=a_1 < \dots < a_k < n} \binom{i}{n-a_k, \dots, a_3-a_2, a_2-i} M_{(n-a_k, \dots, a_2-i, n+1-i)} \end{aligned}$$

For example, $F(L(\Delta^n))^* = \sigma_1^{n+1}$.

Thus the relation between the Ehrenborg F -quasi-symmetric function and the generalized f -polynomial can be illustrated by two commutative diagrams

$$\begin{array}{ccc} \mathcal{P} & \xrightarrow{f} & \text{Qsym}[t_1, t_2, \dots][\alpha] \\ l \downarrow & & \alpha=0 \downarrow \\ \mathcal{R} & \xrightarrow{F^*} & \text{Qsym}[t_1, t_2, \dots] \end{array} \quad \begin{array}{ccc} \mathcal{P} & \xrightarrow{f} & \text{Qsym}[t_1, t_2, \dots][\alpha] \\ L \downarrow & & \Lambda \downarrow \\ \mathcal{R} & \xrightarrow{F^*} & \text{Qsym}[t_1, t_2, \dots] \end{array}$$

For other relations see the next subsection.

14.3 Hopf Comodule Structure

Definition 26. By a *Hopf comodule* (or *Milnor comodule*) over a Hopf algebra X we mean a k -algebra M with a unit provided M is a comodule over X with a coaction $b: M \rightarrow X \otimes M$ such that $b(uv) = b(u)b(v)$, i.e. such that b is a homomorphism of rings.

The ring homomorphism l can be extended to a right graded Hopf comodule structure on $\mathcal{P}$

Proposition 37. *The homomorphism $\Delta: \mathcal{P} \rightarrow \mathcal{P} \otimes \mathcal{R}$:*

$$\Delta(P^n) = \sum_{F \subseteq P^n} F \otimes L(P^n/F) = \sum_{F \subseteq P^n} F \otimes [F, P^n],$$

where by definition $L(P/P) = [P, P]$, defines on $\mathcal{P}$ a right graded Hopf comodule structure over $\mathcal{R}$ such that

$$(\varepsilon \otimes \text{id})\Delta P = 1 \otimes l(P)$$

Here by k we mean $\mathbb{Z}$ or $\mathbb{Q}$. In the latter case we should take $\mathcal{P} \otimes \mathbb{Q}$.

Proof. We should proof that the following three diagrams commute

$$\begin{array}{ccc} \mathcal{P} & \xrightarrow{\Delta} & \mathcal{P} \otimes \mathcal{R} \\ \downarrow \Delta & & \downarrow \text{id} \otimes \Delta \\ \mathcal{P} \otimes \mathcal{R} & \xrightarrow{\Delta \otimes \text{id}} & \mathcal{P} \otimes \mathcal{R} \end{array} \quad \begin{array}{ccc} \mathcal{P} & & \\ \downarrow \Delta & \searrow \text{id} \otimes 1 & \\ \mathcal{P} \otimes \mathcal{R} & \xrightarrow{\text{id} \otimes \varepsilon} & \mathcal{P} \otimes \mathbb{k} \end{array} \quad \begin{array}{ccc} \mathcal{P} & \xrightarrow{l} & \mathcal{R} \\ \downarrow \Delta & & \downarrow 1 \otimes \text{id} \\ \mathcal{P} \otimes \mathcal{R} & \xrightarrow{\varepsilon \otimes \text{id}} & \mathbb{Z} \otimes \mathcal{R} \end{array}$$

and that Δ is a ring homomorphism. Indeed, we have

$$\begin{aligned} (\Delta \otimes \text{id})\Delta P &= (\Delta \otimes \text{id}) \left(\sum_{F \subseteq P} F \otimes [F, P] \right) = \sum_{F \subseteq P} \left(\sum_{G \subseteq F} G \otimes [G, F] \right) \otimes [F, P] = \\ &= \sum_{G \subseteq P} G \otimes \left(\sum_{G \subseteq F \subseteq P} [G, F] \otimes [F, P] \right) = \sum_{G \subseteq P} G \otimes \Delta[G, P] = (\text{id} \otimes \Delta)\Delta P \end{aligned}$$

$$(\text{id} \otimes \varepsilon)\Delta(P) = \sum_{F \subseteq P} F \otimes \varepsilon([F, P]) = P \otimes 1,$$

and

$$(\varepsilon \otimes 1)\Delta P = \sum_{F \subseteq P} \varepsilon(F) \otimes [F, P] = \sum_{v - \text{vertex}} 1 \otimes [v, P] = 1 \otimes \left(\sum_v [v, P] \right) = 1 \otimes l(P).$$

Let us check that Δ is a homomorphism of rings $\mathcal{P} \rightarrow \mathcal{P} \otimes \mathcal{R}$:

$$\begin{aligned} \Delta(PQ) &= \sum_{F \times G \subseteq P \times Q} F \times G \otimes [F \times G, P \times Q] = \sum_{F \subseteq P, G \subseteq Q} F \times G \otimes [F, P] \cdot [G, Q] = \\ &= \left(\sum_{F \subseteq P} F \otimes [F, P] \right) \cdot \left(\sum_{G \subseteq Q} G \otimes [G, Q] \right) = \Delta(P) \cdot \Delta(Q) \end{aligned}$$

$$\Delta(\text{pt}) = \text{pt} \otimes [\text{pt}, \text{pt}] = 1 \otimes 1$$

□

Corollary 38. • Any ring homomorphism $\varphi : \mathcal{P} \rightarrow \mathbb{k}$ defines a ring homomorphism $\mathcal{P} \rightarrow \mathcal{R}$

$$\mathcal{P} \xrightarrow{\Delta} \mathcal{P} \otimes \mathcal{R} \xrightarrow{\varphi \otimes \text{id}} \mathbb{k} \otimes \mathcal{R} \simeq \mathcal{R}$$

• Any homomorphism of abelian groups $\psi : \mathcal{R} \rightarrow \mathbb{Z}$ defines a linear operator $\Psi \in \mathcal{L}(\mathcal{P})$

$$\mathcal{P} \xrightarrow{\Delta} \mathcal{P} \otimes \mathcal{R} \xrightarrow{\text{id} \otimes \psi} \mathcal{P} \otimes \mathbb{Z} \simeq \mathcal{P}$$

• In particular, if $\psi : \mathcal{R} \rightarrow \mathbb{Z}$ is a multiplicative homomorphism, then Ψ is a ring homomorphism.

Example. Let $\varphi = \xi_\alpha$. Then we obtain the ring homomorphism $l_\alpha : \mathcal{P} \rightarrow \mathcal{R}[\alpha]$ defined as

$$l_\alpha(P^n) = (\xi_\alpha \otimes \text{id})\Delta P^n = \sum_{F \subseteq P} \alpha^{\dim F} [F, P]$$

- If we set $\alpha = 0$, then we obtain a usual homomorphism l .
- On the ring of simple polytopes $\mathcal{P}_s$ we have

$$l_\alpha(P^n) = \sum_{F \subseteq P^n} \alpha^{\dim F} \{\hat{0}, \hat{1}\}^{n - \dim F} = \sum_{F \subseteq P^n} \alpha^{\dim F} x^{n - \dim F} = f_1(\alpha, x)$$

is a homogeneous f -polynomial in two variables.

Set $F(\alpha) = \alpha$. Then we have the ring homomorphism $F : \mathcal{R}[\alpha] \rightarrow \text{Qsym}[t_1, t_1, \dots][\alpha]$.

Proposition 39. Let P^n be an n -dimensional polytope. Then

$$F(l_\alpha(P^n)) = f(P^n)^*$$

Proof. The proof is similar to the proof of Proposition 34:

$$F(l_\alpha(P^n)) = \sum_{F^{a_1} \subseteq P} \alpha^{a_1} \sum_{F^{a_1} \subset F^{a_2} \subset \dots \subset F^{a_k} \subset P^n} M_{(a_2 - a_1, \dots, n - a_k)} = \sum_{F^{a_1} \subset \dots \subset F^{a_k} \subset P^n} \alpha^{a_1} M_{(a_2 - a_1, \dots, n - a_k)} = f(P^n)^*$$

□

Proposition 40. The following diagram commutes:

$$\begin{array}{ccc} \mathcal{P} & \xrightarrow{f^*} & \text{Qsym}[t_1, t_2, \dots][\alpha] \\ \Delta \downarrow & & \Delta \downarrow \\ \mathcal{P} \otimes \mathcal{R} & \xrightarrow{f^* \otimes F} & \text{Qsym}[t_1, t_2, \dots][\alpha] \otimes \text{Qsym}[t_1, t_2, \dots] \end{array}$$

Proof. We know that

$$\Delta(M_{(b_1, \dots, b_k)}) = 1 \otimes M_{(b_1, \dots, b_k)} + M_{(b_1)} \otimes M_{(b_2, \dots, b_k)} + \dots + M_{(b_1, \dots, b_{k-1})} \otimes M_{(b_k)} + M_{(b_1, \dots, b_k)} \otimes 1$$

and

$$f(\alpha, t_1, t_2, \dots)(P^n)^* = \alpha^n + \sum_{F^{a_1} \subset \dots \subset F^{a_k}} \alpha^{a_1} M_{(a_2 - a_1, \dots, n - a_k)} = \sum_{F^{a_1} \subset \dots \subset F^{a_k} \subset P^n} \alpha^{a_1} M_{(a_2 - a_1, \dots, n - a_k)}$$

Then

$$\begin{aligned} \Delta(f(P^n)^*) &= \sum_{F^{a_1} \subset \dots \subset G^{a_s} \subset \dots \subset F^{a_k} \subset P^n} \alpha^{a_1} M_{(a_2 - a_1, \dots, a_s - a_{s-1})} \otimes M_{(a_{s+1} - a_s, \dots, n - a_k)} = \\ &= \sum_{G^{a_s} \subseteq P^n} \left(\sum_{F^{a_1} \subset \dots \subset F^{a_{s-1}} \subset G^{a_s}} \alpha^{a_1} M_{(a_2 - a_1, \dots, a_s - a_{s-1})} \right) \otimes \left(\sum_{G^{a_s} \subset F^{a_{s+1}} \subset \dots \subset F^{a_k} \subset P^n} M_{(a_{s+1} - a_s, \dots, n - a_k)} \right) = \\ &= \sum_{G \subseteq P^n} f^*(G) \otimes F([G, P^n]) = (f^* \otimes F) \Delta P^n. \end{aligned}$$

□

These two propositions summarize the relations between the Ehrenborg F -quasi-symmetric function and the generalized f -polynomial.

15 Problem of the Description of Flag Vectors of Polytopes

We have mentioned that on the space of simple polytopes

$$f(\alpha, t_1, t_2, \dots)(P^n) = f_1(\alpha, t_1 + t_2 + \dots)(P^n) = f_1(\alpha, \sigma_1)(P^n).$$

The only linear relation on the polynomial f_1 is $f_1(-\alpha, \alpha + t) = f_1(\alpha, t)$ and it is equivalent to the Dehn-Sommerville relations. In fact, for the polynomial $g = g(\alpha, \sigma_1) \in \text{Qsym}[t_1, t_2, \dots][\alpha]$ this condition is necessary and sufficient to be an image of an integer combination of simple polytopes.

One of the outstanding results in the polytope theory is the so-called g -theorem, which was formulated as a conjecture by P. McMullen [Mc1] in 1970 and proved by R. Stanley [St1] (necessity) and L. Billera and C. Lee [BL] (sufficiency) in 1980.

For an n -dimensional polytope P^n let us define an h -polynomial:

$$h(\alpha, t)(P^n) = h_0 \alpha^n + h_1 \alpha^{n-1} t + \dots + h_{n-1} \alpha t^{n-1} + h_n t^n = f_1(\alpha - t, t)(P^n)$$

Since $f_1(-\alpha, \alpha + t) = f_1(\alpha, t)$, the h -polynomial is symmetric: $h(\alpha, t) = h(t, \alpha)$. So $h_i = h_{n-i}$.

Definition 27. A g -vector is a set of numbers $g_0 = 1$, $g_i = h_i - h_{i-1}$, $1 \leq i \leq \lfloor \frac{n}{2} \rfloor$.

For any positive integers a and i there exists a unique "binomial i -decomposition" of a :

$$a = \binom{a_i}{i} + \binom{a_{i-1}}{i-1} + \dots + \binom{a_j}{j},$$

where $a_i > a_{i-1} > \dots > a_j \geq j \geq 1$.

Let us define

$$a^{(i)} = \binom{a_i + 1}{i + 1} + \binom{a_{i-1} + 1}{i} + \dots + \binom{a_j + 1}{j + 1}, \quad 0^{(i)} = 0.$$

For example, $a = \binom{a}{1}$, so $a^{(1)} = \binom{a+1}{2} = \frac{(a+1)a}{2}$; for $i \geq a$ we have $a = \binom{i}{i} + \binom{i-1}{i-1} + \dots + \binom{i-a+1}{i-a+1}$, so $a^{(i)} = a$.

Theorem (g -theorem, [St1], [BL]). *Integer numbers $(g_0, g_1, \dots, g_{\lfloor \frac{n}{2} \rfloor})$ form a g -vector of some simple n -dimensional polytope if and only if they satisfy the following conditions:*

$$g_0 = 1, \quad 0 \leq g_1, \quad 0 \leq g_{i+1} \leq g_i^{(i)}, \quad i = 1, 2, \dots, \left\lfloor \frac{n}{2} \right\rfloor - 1.$$

As it is mentioned in [Z1] even for 4-dimensional non-simple polytopes the corresponding problem is extremely hard.

Modern tools used to obtain linear and nonlinear inequalities satisfied by flag f -vectors are based on the notion of a cd -index [BK, BE1, BE2, EF, ER, L, St3], a toric h -vector [BE, BL, K, St2, Sts] and its generalizations [F1, F2, L], the ring of chain operators [BH, BLiu, Kalai].

In [St1] R. Stanley constructed for a simple polytope P a projective toric variety X_P so that h_i are the even Betti numbers of the (singular) cohomology of X_P . Then the Poincare duality and the Hard Lefschetz theorem for X_P imply the McMullen conditions for P .

In [Mc2] P. McMullen gave a purely geometric proof of these conditions using the notion of the Polytope algebra.

In [St2] Stanley generalized the definition of the h -vector to an arbitrary polytope P in such a way that if the polytope is rational the generalized $\hat{h}_i$ are the intersection cohomology Betti numbers of the associated toric variety (see the validation of the claim in [Fi]).

The set of numbers $(\hat{h}_0, \dots, \hat{h}_n)$ is called a *toric h -vector*. It is nonnegative and symmetric: $\hat{h}_i = \hat{h}_{n-i}$, and in the case of rational polytope the Hard Lefschetz theorem in the intersection cohomology of the associated toric variety proves the unimodality

$$1 = \hat{h}_0 \leq \hat{h}_1 \leq \dots \leq \hat{h}_{\lfloor \frac{n}{2} \rfloor}$$

Generalizing the geometrical methods of P. McMullen [Mc2] Kalle Karu [K] proved the unimodality for an arbitrary polytope.

The toric h -vector consists of linear combinations of the flag f -numbers. In general case it does not contain the full information about the flag f -vector. In [L] C. Lee introduced an "extended toric" h -vector that carries all the information about the flag f -numbers and consists of a collection of nonnegative symmetric unimodal vectors. In [F1] and [F2] J. Fine introduced the generalized h - and g -vectors that contain toric h - and g -vectors as subsets, but can have negative entries.

Let us look on the problem of the description of flag f -vectors from the point of view of the ring of polytopes.

In the case of simple polytopes we can use the g -theorem to describe all the polynomials $\psi \in \text{Qsym}[t_1, t_2, \dots][\alpha]$ that are images of simple polytopes under the ring homomorphism f :

$$\psi(\alpha, t_1, t_2, \dots) = f(\alpha, t_1, t_2, \dots)(P^n), \quad P^n - \text{a simple } n\text{-dimensional polytope.}$$

In general case Theorem 2 gives a criterion for the polynomial $\psi \in \text{Qsym}[t_1, t_2, \dots][\alpha]$ to be an image of an integer combination of polytopes.

Question 2. Find a criterion for the polynomial $\psi \in \text{Qsym}[t_1, t_2, \dots][\alpha]$ to be an image of an n -dimensional polytope.

References

- [ABS] M. Aguiar, N. Bergeron, F. Sottile, *Combinatorial Hopf Algebras and Generalized Dehn-Sommerville relations*. Compositio Mathematica 142(1) 2006, 1–30, arXiv: math/0310016v1 [math.CO].
- [BR] A. Baker, B. Richter, *Quasisymmetric functions from a topological point of view*. Math. Scand. 103 (2008), 208–242.
- [BB] M. M. Bayer, L. J. Billera, *Generalized Dehn-Sommerville relations for polytopes, spheres and Eulerian partially ordered sets*. Invent. Math. 79: 143–157, 1985.
- [BE] M. M. Bayer and R. Ehrenborg, *The toric h -vectors of partially ordered sets*. Trans. Amer. Math. Soc. 352 (2000), 4515–4531 (electronic).
- [BK] M. M. Bayer, A. Klapper, *A new index for polytopes*. Discrete Comput. Geom. 6, 33–47 (1991).
- [BL] . M. Bayer and C. W. Lee, *Combinatorial aspects of convex polytopes*. In Handbook of convex geometry, Vol. A, B, North-Holland, Amsterdam, 1993, pp. 485–534.

- [BMSW] N. Bergeron, S. Mykytiuk, F. Sottile, and S. van Willigenburg, *Pieri Operations on Posets*. J. of Comb. Theory Series A 91(2000) 84-110. arXiv: math/0002073v2 [math.CO] 21 Jul 2000.
- [BE1] L. Billera, R. Ehrenborg, *Monotonicity of the cd-index for polytopes*. Math. Z. 233(200) 421-441.
- [BE2] L. Billera, R. Ehrenborg and M. Readdy, *The cd-index of zonotopes and arrangements*. Mathematical Essays in Honor of Gian-Carlo Rota, Birkhauser, Boston, 1998.
- [BH] L. Billera, G. Hettley, *Linear Inequalities For Flags In Graded Posets*. J. of Combinatorial Theory Series A, 2000, V89, N1, 77-104, arXiv: math/9706220v1 [math.CO]
- [BL] L. Billera, C. Lee, *A proof of sufficiency of McMullen's conditions for f-vectors of simplicial polytopes*. J. Combin. Theory, Ser. A, 1981, V31, N3, 237-255.
- [BLiu] L. Billera, N. Liu, *Non-commutative enumeration in graded posets*. Journal of Algebraic Combinatorics 12 (2000), 7-24.
- [BS] R. Bott, H. Samelson, *On the Pontryagin product in spaces of paths*. Comment. Math. Helv., 27, 1953, 320-337
- [BR] V. M. Buchstaber, N. Ray, *An invitation to toric topology: Vertex four of a remarkable tetrahedron*. Toric topology. Contemporary Mathematics (AMS), v. 460, Providence, R.I., 2008, 1-27; MIMS, 2008.31: <http://www.manchester.ac.uk/mims/eprints>.
- [Buch] V. M. Buchstaber, *Ring of Simple Polytopes and Differential Equations*. Proceedings of the Steklov Institute of Mathematics, v. 263, 2008, 1-25.
- [CFL] K. T. Chen, R. H. Fox, and R. C. Lyndon, *Free differetional calculus*. IV, Ann. Math. 68 (1958), 81-95.
- [Ehr] R. Ehrenborg, *On Posets and Hopf Algebras*. Advances in Mathematics, 119, 1-25 (1996).
- [EF] R. Ehrenborg, H. Fox, *Inequalities for cd-indices of joins and products of polytopes*. Combinatorica 23 (2003), 427-452.
- [ER] R. Ehrenborg, M. Readdy, *Coproducts and the cd-index*. J. Algebraic Combin. 8 (1998), 273-299.
- [Fi] K. -H. Fieseler, *Rational Intersection Cohomology of Projective Toric Varieties*. Journal f.d. reine u. angew. Mathematik 413 (1991), 88-98.
- [F1] J. Fine, *A complete h-vector for convex polytopes*. preprint, arXiv:0911.5722v1 [math.CO]
- [F2] J. Fine, *A complete g-vector for convex polytopes*. preprint, arXiv:1001.1562v1 [math.CO]
- [Fr] A. Frohmader, *Face vectors of flag complexes*. Israel Journal of Mathematics, 164, 2008, 153-164.
- [Gb] B. Grunbaum, *Convex Polytopes*. vol. 221 of Graduate Texts in Mathematics, Springer-Verlag, New York, Second ed., 2003.
- [H] M. Hazewinkel, *The Algebra of Quasi-Symmetric Functions is free over integers*. Advances in Mathematics 164, 283-300 (2001).
- [JR] S. Joni and G.-C. Rota, *Coalgebras and bialgebras in combinatorics*. Stud. Appl. Math. 61(1979), 93-139.
- [Kalai] G. Kalai, *A new basis for polytopes*. J. Combinatorial Theory, Ser A 49 (1988), 191-208.
- [K] Kalle Karu, *Hard Lefschetz Theorem for Nonrational Polytopes*. Invent. Math. 157 (2004), 419-447, arXiv: math/0112087v4 [math.AG] 17 Dec 2002.
- [L] Carl W. Lee, *Sweeping the cd-Index and the Toric h-Vector*. 2009, preprint: <http://www.ms.uky.edu/~lee/cd.pdf>
- [Mc1] P. McMullen, *The numbers of faces of simplicial polytopes*. Israel J.Math. 1971, V9, 559-570.

- [Mc2] P. McMullen, *On Simple Polytopes*. Invent. math. 113, 419-444 (1993).
- [N] S. P. Novikov, *Various doublings of Hopf algebras. Operator algebras on quantum groups, complex cobordisms*. Russian Mathematical Surveys, 47:5, 1992, 198–199.
- [RS] N. Ray, W. Schmitt, *Combinatorial models for coalgebraic structures*. Advances in Mathematics 1998, Vol 138, Iss 2, 211-262.
- [Re] C. Reutenauer, *Free Lie algebras*. Oxford Univ. Press, Oxford, UK, 1993.
- [Sch1] William Schmitt, *Antipodes and incidence coalgebras*. Journal of Combinatorial Theory A, 46:264-290, 1987.
- [Sch2] William Shcmitt, *Incidence Hopf Algebras*. J.Pure Appl. Algebra 96(1994), 299-330.
- [St1] R. P. Stanley, *The number of faces of simplitial convex polytope*. Advances in Math. 1980. V. 35, N3, 236-238.
- [St2] R. P. Stanley, *Generalized h-vectors, intersection cohomology of toric varieties, and related results*. Commutative Algebra and Combinatorics, Advanced Studies in Pure Mathematics 11, Kinokuniya, Tokyo, and North-Holland, Amsterdam/New York, 1987, pp. 187-213.
- [St3] R. P. Stanley, *Flag f-vectors and the cd-index*. Math. Z. 216, 483–499 (1994).
- [Sts] C. Stenson, *Relationships among flag f-vector inequalities for polytopes*. Discrete and Computational Geometry 31 (2004), pp. 257-273
- [Z1] G. M. Ziegler, *Face numbers of 4-polytopes and 3-spheres*. Proceedings of the ICM 2002 Beijing, Volume III, 2004, 625–634; arXiv: math/0208073v2 [math.MG] 20 Jun 2003.
- [Z2] G. M. Ziegler, *Lectures on Polytopes*. Springer-Verlag, New York, 2007.